

518,966

Rec'd PCT

21 DEC 2004

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
31 December 2003 (31.12.2003)


PCT

(10) International Publication Number  
**WO 2004/000993 A2**

- (51) International Patent Classification<sup>7</sup>: C12N
- (21) International Application Number: PCT/CA2003/000897
- (22) International Filing Date: 13 June 2003 (13.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 2,391,118 21 June 2002 (21.06.2002) CA
- (71) Applicants (for all designated States except US): UNIVERSITE DU QUEBEC A MONTREAL [CA/CA]; Succursale Centre-Ville, Case Postale 8888, Montreal, Quebec H3C 3P8 (CA). CENTRE HOSPITALIER DE L'UNIVERSITE DE MONTREAL [CA/CA]; Centre de Recherche Hotel Dieu, Pavillon Masson, 3850 Rue St-Urbain, Montreal, Quebec H2W 1T7 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): PAQUIN, Joanne [CA/CA]; 10151 Georges-Baril, Montréal, Québec H2C 2M9 (CA). GUTKOWSKA, Jolanta [CA/CA]; 139 Avenue Appin, Ville Mont-Royal, Québec H3P 1V6 (CA). DANALACHE, Bogdan, A. [RO/CA]; 4555 St. Kevin, App. 306, Montréal, Québec H3T 1J1 (CA). JANKOWSKI, Marek [CA/CA]; 5155 West Broadway, Montreal, Quebec H4V 2A1 (CA).
- (74) Agents: SHAHINIAN, Serge, S. et al.; SMART & BIGGAR, 1000 de la Gauchetiere Street West, Suite 3400, Montréal, Québec H3B 4W5 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,



[Continued on next page]

(54) Title: OXYTOCIN AS CARDIOMYOGENESIS INDUCER AND USES THEREOF




P19 CELLS

DAYS  
0  
2  
4  
14



DMSO-INDUCED CELLS



OT-INDUCED CELLS

(57) Abstract: The invention relates to oxytocin and oxytocin-related compounds and functional derivatives thereof, and uses thereof to induce differentiation of a non-cardiomyocyte (e.g. a stem/progenitor cell) to a cardiomyocyte. The invention further relates to the methods of prevention or treatment of conditions characterized by or associated with a cardiomyocyte loss or deficiency, by administration of oxytocin or an oxytocin-related compound or a functional derivative thereof to a subject, or via the administration/transplantation of a cell differentiated ex vivo by a method of the invention. The invention further relates to methods, uses, commercial packages and culture media relating to such differentiation and prevention/treatment.

WO 2004/000993 A2

BEST AVAILABLE COPY



SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**OXYTOCIN AS CARDIOMYOGENESIS INDUCER  
AND USES THEREOF**

**5 FIELD OF THE INVENTION**

The present invention relates to the field of cell differentiation. More particularly, the present invention relates to the use of oxytocin (OT) as a cell differentiating agent, and even more specifically as an inducer of cardiomyogenesis. The present invention further relates to the use of cardiomyocytes obtained by oxytocin-induced differentiation of non-cardiomyocytes, in the treatment of diseases associated with loss of cardiomyocytes, such as congenital and aging-related heart pathologies.

**BACKGROUND OF THE INVENTION**

Each year, up to 7% of the three million newborns in the USA have birth defects, many of which predominantly affect the heart. Furthermore, the great prevalence of cardiovascular diseases in aging populations is well known. In Canada and the US, these diseases account for about 45% of all deaths. A number of drugs exist for the treatment of such diseases. Some drugs are used to improve the cardiodynamic properties of the heart (e.g. agonists/antagonists of adrenergic receptors), while others are used to reduce prejudicing conditions to disease (e.g. substances that attenuate hypercholesterolemia). In some cases, the cardiovascular diseases are treated by surgical intervention.

Today, new prospective therapies envisage myocardial regeneration as an alternative for treating cardiovascular diseases because heart infarction, congestive heart failure and acute myocardial ischemia lead to an irreversible death of cardiac tissue (cardiomyocytes and vascular structures) which

becomes replaced by scar tissue. However, there is currently no established cardiac regenerative therapy.

Therefore, there is a need for new prospective therapies and new drugs to prevent and treat heart-related diseases.

5

#### SUMMARY OF THE INVENTION

The present invention pertains to the use of oxytocin  
10 (OT), functional derivatives, and/or physiological precursors thereof, and nucleic acids capable of encoding OT, its derivatives and/or precursors noted above, as a cell differentiating agent and in compositions useful for treating or preventing diseases, such as heart diseases and in  
15 particular those associated with loss of cardiomyocytes. More particularly, the present invention pertains to the use of oxytocin or an oxytocin-related compound, corresponding gene construct(s) and/or their functional derivatives as an inducer of cardiomyogenesis, and more specifically as an inducer that  
20 promotes the differentiation of non-cardiomyocytes (e.g. stem/progenitor cells) *in situ*, which can be used to repair, restore or fortify damaged or otherwise inadequate cardiac tissue. The present invention also pertains to the use of oxytocin and functional derivatives thereof to induce cardiac  
25 differentiation of non-cardiomyocytes (e.g. stem/progenitor cell) in cell culture in order to provide material for cell or tissue grafting in the heart.

According to a first aspect, the invention provides a method of inducing differentiation of a non-cardiomyocyte into  
30 a cardiomyocyte, said method comprising stimulating oxytocin receptor (OTR) activity in said non-cardiomyocyte. In an embodiment, the method comprises contacting said non-cardiomyocyte with an agent capable of stimulating OTR activity.

35 In an embodiment, the above-mentioned method is performed

*in vitro*. In an embodiment, the above-mentioned method is performed *in vivo*.

The invention further provides a method of treating a disease characterized by cardiomyocyte loss or deficiency in an animal, said method comprising stimulating oxytocin receptor (OTR) activity in a non-cardiomyocyte cell of said animal. In an embodiment, the method comprises administering an agent capable of stimulating OTR activity to said animal. In an embodiment, the method comprises administering a nucleic acid capable of encoding oxytocin or a functional derivative thereof to said animal.

The invention further provides a method of treating a disease characterized by cardiomyocyte loss or deficiency in an animal, said method comprising: inducing, using the above-mentioned method, differentiation of a non-cardiomyocyte cell into a cardiomyocyte; and implanting said cardiomyocyte into said animal. In an embodiment, the method comprises contacting the non-cardiomyocyte with an agent capable of stimulating OTR activity. In an embodiment, the method comprises introducing into the non-cardiomyocyte a nucleic acid capable of encoding oxytocin or a functional derivative thereof. In an embodiment, the non-cardiomyocyte is autologous to said animal. In an embodiment, the method further comprises obtaining said non-cardiomyocyte from said animal prior to inducing said differentiation. In an embodiment, the non-cardiomyocyte is non-autologous to said animal, in a further embodiment, allogenic to said animal, in yet a further embodiment, xenogenic to said animal.

The invention further provides a use of an agent capable of stimulating OTR activity for treating a disease characterized by cardiomyocyte loss or deficiency in an animal.

The invention further provides a commercial package comprising an agent capable of stimulating OTR activity together with instructions for treating a disease characterized by cardiomyocyte loss or deficiency in an animal.

The invention further provides a commercial package comprising an agent capable of stimulating OTR activity together with instructions for inducing differentiation of a non-cardiomyocyte to a cardiomyocyte.

5 The invention further provides a commercial package comprising a culture medium comprising oxytocin or a functional derivative thereof; together with instructions for culturing a non-cardiomyocyte in said culture medium thereby to induce differentiation of said non-cardiomyocyte into a cardiomyocyte.

10 In an embodiment, the non-cardiomyocyte is a mammalian non-cardiomyocyte, in a further embodiment, a human non-cardiomyocyte. In an embodiment, the oxytocin or functional derivative thereof is present in said medium at a concentration from about  $10^{-10}$  M to about  $10^{-4}$  M, in a further embodiment,  
15 from about  $10^{-9}$  M to about  $10^{-6}$  M, in yet a further embodiment, from about  $10^{-8}$  M to about  $10^{-7}$  M.

The invention further provides a composition for treatment of a disease characterized by cardiomyocyte loss or deficiency comprising oxytocin or a functional derivative  
20 thereof in and a pharmaceutically acceptable carrier.

The invention further provides a method of identifying or characterizing a compound for inducing differentiation of a non-cardiomyocyte into a cardiomyocyte, said method comprising:  
25 (a) contacting a test compound with a cell comprising an oxytocin receptor (OTR) or an OTR-encoding nucleic acid; and  
(b) determining whether OTR activity or expression is increased in the presence of the test compound, said increase in OTR activity or expression being an indication that the test compound may be used for inducing differentiation of a non-  
30 cardiomyocyte into a cardiomyocyte.

The invention further provides a method of identifying or characterizing a compound for treatment of a disease characterized by cardiomyocyte loss or deficiency, said method comprising: (a) contacting a test compound with a cell  
35 comprising an oxytocin receptor (OTR) or an OTR-encoding

nucleic acid; and (b) determining whether OTR activity or expression is increased in the presence of the test compound, said increase in OTR activity or expression being an indication that the test compound may be used for for treatment of a disease characterized by cardiomyocyte loss or deficiency.


The invention further provides a method of identifying a cell capable of differentiation to a cardiomyocyte, said method comprising determining whether OTR activity or expression is present in said cell, said presence being an indication that said cell is capable of differentiation to a cardiomyocyte.

The invention further provides a commercial package comprising means for determining OTR activity or expression together with instructions for identifying a cell capable of differentiation to a cardiomyocyte.

In an embodiment, the above-mentioned agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).

In an embodiment, the above-mentioned oxytocin or functional derivative thereof has the structure:

20



wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.

In an embodiment, the above-mentioned method comprises introducing into the non-cardiomyocyte a nucleic acid capable of encoding oxytocin or an oxytocin-related compound. In an embodiment, the nucleic acid is selected from the group consisting of: (a) SEQ ID NO:5; (b) a nucleic acid sequence capable of encoding SEQ ID NO:6; and (c) a nucleic acid sequence substantially identical to (a) or (b).

In an embodiment, the non-cardiomyocyte is a mammalian non-cardiomyocyte, in a further embodiment, a human non-cardiomyocyte.

In an embodiment, the non-cardiomyocyte is a stem or progenitor cell. In an embodiment, the stem or progenitor cell is selected from the group consisting of embryonic and adult stem or progenitor cells. In an embodiment, the stem or progenitor cell is selected from the group consisting of circulating and non-circulating stem or progenitor cells.

In an embodiment, the above-mentioned cardiomyocyte is characterized by an alteration of a phenotypic feature relative to said non-cardiomyocyte, wherein said phenotypic feature is selected from the group consisting of: (a) level of oxytocin receptor (OTR) protein or OTR-encoding nucleic acid; (b) level of ANP protein or ANP-encoding nucleic acid; (c) level of muscular MHC protein or MHC-encoding nucleic acid; (d) level of DHPR-alpha1 protein or DHPR-alpha1-encoding nucleic acid; (e) level of sarcomeric marker proteins; (f) level of ion channels; (g) mitochondrial dye retention; (h) appearance of rhythmic beats; and (i) chronotropic responses.

In an embodiment, the above-mentioned animal is a mammal, in a further embodiment, a human.

In an embodiment, the above-mentioned disease is selected from the group consisting of cardiac congenital dysfunctions, aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial ischemia.

The invention further provides a pharmaceutical composition which comprises oxytocin and/or of a functional derivative of oxytocin in an amount effective to promote and/or induce differentiation of a non-cardiomyocyte (e.g. stem/progenitor cell) into a cardiomyocyte, and a suitable pharmaceutical acceptable diluent or carrier.

According to another aspect of the invention, oxytocin and/or its functional derivatives, are used as an active agent in the preparation of a medication for preventing or treating a heart disease or for treating an injury to cardiac tissues. The invention also provides methods for preventing or treating a heart disease or for treating an injury to cardiac tissues,



comprising the administration to a patient in need thereof of a therapeutically effective amount of oxytocin or of a functional derivative of oxytocin or the administration of a therapeutically effective amount of a composition as defined  
5 hereinabove.

According to a further aspect, the invention provides a method for inducing and/or promoting differentiation of cells and more particularly stem/progenitor cells cultured *in vitro* into cardiac cells, such as cardiomyocytes. In a preferred  
10 embodiment, the method comprises the step of providing to the *in vitro* cultured stem/progenitor cells an effective amount of oxytocin or of a functional derivative thereof. According to another aspect, the present invention provides a method to stimulate the fusion of newly-differentiated cardiomyocytes.

15 An advantage of the present invention is that it provides effective means for maintaining or stimulating the regeneration of cardiac cells, such as cardiomyocytes, and thereby, it permits the treatment of injuries to the heart tissues. Another advantage of the present invention is that it improves the  
20 efficiency of methods for culturing cardiac cells *in vitro* either as model system or graft material.

Other objects and advantages of the present invention will be apparent upon reading the following non-restrictive description of several preferred embodiments made with  
25 reference to the accompanying drawings.

#### BRIEF DESCRIPTION OF THE DRAWINGS

30 **Figure 1** is a diagram showing the time schedule of the differentiation of P19 cells to cardiomyocytes. P19 cells were cultivated as aggregates from day 0 to day 4 in the presence of DMSO (0.5% w/v) or oxytocin (OT) ( $10^{-7}$  M) as the agent inducing cellular differentiation. At day 4, aggregates

(embryoid bodies) were transferred to tissue culture dishes or multiwell plates and grown in the absence of the agent. Micrographs (100X magnification) show undifferentiated cells and day 14 cardiomyocyte derivatives obtained after DMSO or OT treatment.

**Figures 2A and 2B** show that oxytocin (OT) induces myocyte immunological markers in P19 cells. P19 cell aggregates were treated from day\_0 to day 4 with DMSO, OT or no differentiation agent, and stained on day 14 with anti-MHC or anti-DHPR-alpha1 antibodies. **Figure 2A** are micrographs (100X magnification) showing day 14 cells that were exposed to OT treatment. Normal light and fluorescence pictures are presented side by side. **Figure 2B** is a graph showing immunoreactivity (ir) signals obtained for undifferentiated cells grown in monolayers (Undiff.), non-treated cell aggregates (No inducer) and cell aggregates treated with DMSO or OT. Immunoreactive foci were absent (0), very rare (slightly above zero), or abundant (++ and +++). Results are representative of 3 independent experiments. Although not presented, aggregates were also treated for 6 days with OT. There was no difference with the 4-day treatment.

**Figures 3A, 3B and 3C** show comparison of the cardiomyogenic effect of oxytocin (OT) and DMSO. **Figure 3A** shows the retention of rhodamine<sup>123</sup> in non-induced and induced P19 cultures. P19 cells were cultured as aggregates for 4 days in the absence (No inducer) or the presence of OT or DMSO, using 1 petri dish per treatment. At day 4, aggregates of each petri dish were evenly distributed in wells of a 24-well tissue culture plate. At day 8, the cells were incubated for 45 min in the presence of 1 µg/ml of the dye, washed extensively, and cultured in complete medium without dye for 48 h. The photograph shows rhodamine<sup>123</sup> retention by cells induced by OT at day 10 of culture. The

retained dye was fluorimetrically quantified for each well, and the results are reported as the means  $\pm$  SEM of 24 determinations. The symbol \* indicates a highly significant difference with No inducer, and symbol #, a highly significant difference between OT and DMSO treatments ( $p < 0.001$ ). **Figure 3B** is a graph showing the time course of appearance of beating cell colonies upon treatment with different agents. Aggregates of 1 petri dish treated for 4 days with the indicated agent(s) were evenly distributed in wells of a 24-well tissue culture plate. Then, each plate was examined at 2-day intervals for the number of wells containing beating cell colonies. The results are representative of 3 independent differentiation experiments. **Figure 3C** shows the RT-PCR analysis of ANP gene transcript in undifferentiated and induced cultures. Cell aggregates were exposed to OT or DMSO in the absence or presence of OTA from day 0 to day 4, and RNA was extracted at day 14 of the differentiation protocol. ANP transcript was also evaluated in undifferentiated cells grown in monolayers (Undiff.). Mouse heart ventricle mRNA was used as a positive control. Levels of ANP mRNA were adjusted by dividing by corresponding GAPDH mRNA and then expressed as the percentage of the Undiff. value. Results are reported as the means  $\pm$  SEM of 5 independent studies. The symbol \* indicates a significant difference with Undiff., and symbol \$, a significant difference between OT and OT + OTA treatments ( $p < 0.05$ ).

**Figures 4A, 4B, and 4C** show that OT and DMSO increase OTR expression in P19 cells. P19 cells were cultured as aggregates for 4 days in the absence (No inducer) or presence of DMSO (0.5%), OT ( $10^{-7}$  M) and/or OTA ( $10^{-7}$  M), and then plated in tissue culture dishes where they grew in the absence of the agent. At day 14 of differentiation, the cells were examined for OTR expression, together with undifferentiated (Undiff.) cells grown in monolayers. The results are representative of

3 independent differentiation experiments. **Figure 4A** are micrographs showing the immunocytochemistry results. **Figure 4B** shows the immunoblotting results (20 µg protein/lane). **Figure 4C** shows the RT-PCR analysis.

5

**Figures 5A and 5B** show results of studies of the concentration dependence of the cardiomyogenic effect of OT. P19 were induced with different concentrations of OT or with DMSO for 4 days and cultured in absence of inducer for 10 more days. (A) Wells were scored for the presence of beating cell colonies. (B) OTR mRNA expression was analyzed on day 14. Results were compared to non-induced (NI) cells.

**Figure 6** shows that P19 cells induced with OT and DMSO produce OT. P19 were induced with OT or DMSO for 4 days and cultured in absence of inducer for 10 more days. Cells were incubated for 4 h in serum-free medium and collected with their culture media for analysis of OT production by RIA.

**Figure 7** shows that OT/OTR system is involved in RA-induced cardiac differentiation of P19 cells. P19 cells were induced with OT ( $10^{-7}$  M), RA ( $10^{-7}$  M) or their combination for 4 days, in the absence or presence of OTA ( $10^{-7}$  M). Culture was pursued for 10 more days in absence of inducer. Wells were scored for the presence of beating cell colonies throughout (A), and OTR mRNA expression analyzed on day 14 (B).

**Figure 8** shows results of studies of morphology of cells at day 6 of differentiation. P19 cells were induced with  $10^{-8}$  M RA (a cardiomyogenic concentration) in the absence or presence of  $10^{-7}$  M OTA. Morphology was examined two days after the removal of inducer. Cell populations treated with RA + OTA contained some neurons as indicated by the appearance of neuritic processes.

**Figure 9** shows results demonstrating that OT antagonist (OTA) completely inhibits P19 cell differentiation into cardiomyocytes as induced by  $T_3$ , as well as by DMSO, retinoic acid (RA). P19 cells were treated for 4 days with 30 nM  $T_3$ , 0.5% DMSO or  $10^{-7}$  M RA, in the absence or presence of  $10^{-7}$  M OTA. Wells were then scored for the presence of beating cell colonies.

**Figure 10** shows results of studies of the *in vivo* effect of retinoic acid (RA) in the fetal heart (E21). (A) OT concentration measured by radioimmunoassay. (B) OT mRNA as shown by semiquantitative RT-PCR. (C)  $^{125}\text{I}$ -OTA binding to the fetal sections demonstrated by autoradiography. The representative competition curve of  $^{125}\text{I}$ -OTA binding to the fetal heart sections by unlabelled OTA. (D) Retinal dehydrogenase 1 transcript detected by Northern blotting in the fetal heart (1) and fetal kidney (2).

**Figure 11** sets forth the DNA sequence (SEQ ID NO:5) of the human oxytocin gene and the encoded polypeptide (SEQ ID NO:6). The signal peptide is underlined. The oxytocin sequence is shown in bold. The remaining C-terminal portion corresponds to neurophysin I. The nucleotide sequence and peptide sequences corresponding to the mature OT peptide are further shown.

**Figure 12** sets forth the DNA sequence (SEQ ID NO:7) of the human oxytocin receptor gene and the encoded polypeptide (SEQ ID NO:8).

**DETAILED DESCRIPTION OF THE INVENTION**

The present invention generally pertains to the use of oxytocin or an oxytocin-related compound, corresponding gene construct(s) and/or their functional derivative(s) as a cell-differentiating agent, which may in embodiments be used in compositions useful for treating or preventing heart diseases, in particular those associated with loss of cardiomyocytes. More particularly, the present invention pertains to the use of oxytocin, its gene construct and/or functional derivatives thereof as an inducer of cardiomyogenesis, and more specifically as an inducer that promotes heart regeneration via the differentiation of non-cardiomyocytes into cardiomyocytes (e.g. stem/progenitor cells) *in situ*. The present invention also pertains to the use of oxytocin or an oxytocin-related compound, corresponding gene construct(s) and/or their functional derivative(s) to induce cardiac differentiation of a non-cardiomyocyte (e.g. a stem/progenitor cell) in cell culture in order to provide material for cell or tissue grafting in the heart. As used herein, the term "non-cardiomyocyte" refers to any cell that lacks at least one phenotypic feature typical of a cardiomyocyte, and is capable of differentiation to a cardiomyocyte. As used herein, the term "stem/progenitor cell" refers to any cell having the capacity of being differentiated into a cell with altered or additional functional characteristics, such as a cardiomyocyte. Stem/progenitor cells include pluripotent stem cells capable of differentiation into a variety of cell types. Preferred stem/progenitor cells contemplated by the present invention are embryonic stem cells, stem cells of developed tissues, and cells of a developed phenotype but still capable of transdifferentiation, i.e. to differentiate to another cell phenotype.

It is demonstrated herein that various agents (e.g. oxytocin, retinoic acid and triiodothyronine) may be used to

differentiate a non-cardiomyocyte into a cardiomyocyte. The differentiating activity of these agents is shown to act via stimulation of the oxytocin receptor (OTR), as such differentiation is inhibited in the presense of the OTR antagonist OTA. Thus, it is demonstrated herein that stimulation of OTR activity in a non cardiomyocyte may be used to differentiate the non-cardiomyocyte to a cardiomyocyte. Accordingly, the invention provides a method of differentiating or inducing differentiation of a non-cardiomyocyte to a cardiomyocyte, the method comprising stimulating or inducing OTR activity in the non-cardiomyocyte. In an embodiment, the method comprises contacting the non-cardiomyocyte with an agent capable of stimulating or inducing OTR activity. "An agent capable of stimulating/inducing OTR activity" as used herein refers to any agent which when introduced to or contacted with an appropriate system or cell, results in a stimulation of OTR activity in that system or cell. In embodiments, such agents include OTR agonists (e.g. oxytocin or related compounds or functional derivatives thereof) as well as other varieties of agents capable of such stimulation, such as retinoic acid and triiodothyronine.

The present invention thus provides a novel cell differentiating agent and more particularly a new cardiomyogenic factor. As used herein, "cardiomyogenic factor" refers to any compound (or to any mixture of compounds) that promotes the genesis, maturation, growth, and regeneration of cardiac cells, and more specifically promotes differentiation of a non-cardiomyocyte (e.g. a stem/progenitor cell) into a cardiomyocyte.

A cardiomyocyte may be distinguished from a non-cardiomyocyte based on an alteration of a variety of phenotypic features, including, but not limited to the level of oxytocin receptor (OTR) protein or OTR-encoding nucleic acid; level of ANP protein or ANP-encoding nucleic acid; level of muscular/cardiac myosin heavy chain (MHC) protein or MHC-

encoding nucleic acid; level of dihydropyridine receptor-alpha1 (DHPR $\alpha$ 1) protein or DHPR $\alpha$ 1-encoding nucleic acid; level of sarcomeric marker proteins; level of ion channels; mitochondrial dye retention; appearance of rhythmic beats and  
5 chronotropic responses.


In an embodiment, the present invention describes the use of oxytocin in a pharmaceutical composition and in a method for promoting the genesis, maturation, growth, and regeneration of cardiac cells. The cardiac cells that are most susceptible to  
10 benefit from the composition of the invention are stem cells and newly differentiated cardiomyocytes. Also, the present invention relates to the use of oxytocin for the preparation of a composition or a medicament for the treatment or prevention of heart diseases and in particular those associated  
15 with loss of cardiomyocytes.

The pharmaceutical composition of the invention thus comprises oxytocin and/or of a functional derivative oxytocin in an amount effective to promote and/or induce differentiation of stem/progenitor cells into cardiac cells, and a suitable  
20 pharmaceutical acceptable diluent or carrier.

Oxytocin is a nonapeptide with two cysteine residues that form a disulfide bridge between positions 1 and 6 and corresponds to the formula:




The non-(carboxy-terminal) amidated version has the following structure:



35 Another version of OT retains an additional C-terminal Gly



residue (retained from the proprotein), and has the following structure.



Cys-Tyr-Ile-Gln-Asn-Cys-Pro-Leu-Gly-Gly (SEQ ID NO:15)

Thus, the oxytocin or an oxytocin-related compound, and/or their functional derivative(s) according to the present invention are in embodiments substantially pure oxytocin produced by chemical synthesis, or purified from plasma and various tissues, but preferably from the pituitary gland, or produced by recombinant techniques. As generally understood and used herein, the term substantially pure refers to an oxytocin preparation that is generally lacking cellular or other undesirable components.

"oxytocin-related compound" or "oxytocin-like compound" refers to a compound which is structurally and/or functionally related to oxytocin. Such compounds include homologs, variants or fragments of oxytocin which retain oxytocin activity. Such compounds may comprise a peptide which is substantially identical to oxytocin or fragment thereof, e.g. substantially identical to the human oxytocin peptide set forth above and in Figure 11. Similarly, such compounds include peptides or proteins encoded by a nucleic acid sequence which is substantially identical to, or is related by hybridization criteria (see below) to a nucleic acid sequence capable of encoding oxytocin, such as the region of human oxytocin DNA set forth in Figure 11. Such compounds further include precursors (e.g. naturally-occurring precursors) or prodrugs which are metabolized or otherwise converted to an active compound at the site of action.

A "functional derivative", as is generally understood and used herein, refers to a polypeptide sequence that possesses a functional biological activity that is substantially similar

to the biological activity of the whole protein sequence. A functional derivative of a protein may or may not contain post-translational modifications such as covalently linked carbohydrate, or disulphide bonds, if such modification is not  
5 necessary for the performance of a specific function. The term "functional derivative" is intended to the "fragments", "segments", "variants", "analogs" or "chemical derivatives" of a protein.

Oxytocin-related compounds, also include precursors  
10 which can be modified to produce mature, active oxytocin, or an analog having oxytocin activity. In this regard, the human oxytocin gene encodes two proteins, oxytocin and neurophysin I. Oxytocin is therefore naturally produced as a proprotein of oxytocin-neurophysin I, as shown in Figure 11 (human  
15 oxytocin gene DNA [SEQ ID NO:3] and encoded polypeptide [SEQ ID NO:4]; Genbank accession NM\_00915.2), which is subsequently modified to the active oxytocin peptide. With reference to the DNA sequence in Figure 11, the regions of the sequence correspond to the following:

		<u>DNA region</u>
20	coding sequence	nucleotides 37-414
	signal peptide	37-93
	proprotein	94-411
	mature oxytocin peptide	94-120
	mature neurophysin I peptide	130-408

25 With reference to the polypeptide sequence of Figure 11, the signal peptide is underlined and the sequence of oxytocin is shown in bold.

The terms "fragment" and "segment" as are generally understood and used herein, refer to a section of a protein,  
30 and are meant to refer to any portion of the amino acid sequence.

The term "variant" as is generally understood and used herein, refers to a protein that is substantially similar in structure and biological activity to either the protein or  
35 fragment thereof. Thus two proteins are considered variants if

they possess a common activity and may substitute each other, even if the amino acid sequence, the secondary, tertiary, or quaternary structure of one of the proteins is not identical to that found in the other.

5       The term "analog" as is generally understood and used herein, refers to a protein that is substantially similar in function to oxytocin. Preferred OT analogs include for instance extended forms of OT such as OT-Gly, OT-Gly-Lys and OT-Gly-Lys-Arg. These extended forms are biological oxytocin precursors  
10 in vivo.

As used herein, a protein is said to be a "chemical derivative" of another protein when it contains additional chemical moieties not normally part of the protein, said moieties being added by using techniques well known in the art.  
15 Such moieties may improve the protein solubility, absorption, bioavailability, biological half life, and the like. Any undesirable toxicity and side effects of the protein may be attenuated and even eliminated by using such moieties. For example, OT and OT fragments can be covalently coupled to  
20 biocompatible polymers (polyvinyl-alcohol, polyethylene-glycol, etc) in order to improve stability or to decrease antigenicity.

The amount of oxytocin and/or functional derivatives thereof present in the composition of the present invention is a therapeutically effective amount. A therapeutically effective  
25 amount of oxytocin is that amount of oxytocin or derivative thereof necessary so that the protein acts as a cardiomyogenic factor, and more particularly the amount necessary so that the protein promote the generation, maturation, growth, and restoration of cardiac cells, and more specifically,  
30 cardiomyocytes, and the fortification of cardiac tissue with such cells. The exact amount of oxytocin and/or functional derivatives thereof to be used will vary according to factors such as the protein biological activity, the type of condition being treated as well as the other ingredients in the  
35 composition. Typically, the amount of oxytocin should vary from

about  $10^{-15}$  M to about  $10^{-2}$  M. In a preferred embodiment, oxytocin is present in the composition in an amount from about  $10^{-10}$  M to about  $10^{-4}$  M, preferably from about  $10^{-9}$  M to about  $10^{-6}$  M. In the embodiments, the composition comprises about  $10^{-7}$  of oxytocin for *in vivo* applications and  $10^{-6}$  M of oxytocin for *ex vivo* applications.

A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result, such as reduction of progression of a disease characterized by cardiomyocyte loss or deficiency. A therapeutically effective amount of oxytocin, or an oxytocin-related compound, may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the compound to elicit a desired response in the individual. Dosage regimens may be adjusted to provide the optimum therapeutic response. A therapeutically effective amount is also one in which any toxic or detrimental effects of the compound are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result, such as preventing or inhibiting onset or progression of a disease characterized by cardiomyocyte loss or deficiency. A prophylactically effective amount can be determined as described above for the therapeutically effective amount. For any particular subject, specific dosage regimens may be adjusted over time according to the individual need and the professional judgement of the person administering or supervising the administration of the compositions.

Further therapeutic agents can be added to the composition of the invention. For instance, the composition of the invention may also comprise therapeutic agents such as modulators of heart function (agonists/antagonists of adrenergic receptors, activators of neurohormones, cytokines, signaling second messengers such as cAMP / cGMP / calcium or

their analogs, inhibitors of the degradation of second messengers); growth factors, steroid / glucocorticoid / retinoid / thyroid hormones which modulate heart gene expression; proteases / protease inhibitors / cell adhesion proteins / angiogenic factors that modulate cardiac tissue organization and/or vascularization; antioxidants that provide cell protection to endogenous cardiac tissue as well as to exogenous cardiomyocyte cultures before, during and after engrafting; anticoagulants; immunosuppressive drugs.

Further to the therapeutic agents, the pharmaceutical compositions of the invention may also contain metal chelators (proteinic or not), metal scavengers (proteinic or not), coating agents, preserving agents, solubilizing agents, stabilizing agents, wetting agents, emulsifiers, sweeteners, colorants, odorants, salts, buffers, coating agents and/or antioxidants. For preparing such pharmaceutical compositions, methods well known in the art may be used.

The method of preparation of the composition of the invention consists simply in the mixing of purified oxytocin and other component(s) in a suitable solution in order to get a homogenous physiological suspension. A suitable solution is an isotonic buffered saline solution comprising sodium, potassium, magnesium or manganese, and calcium ions at physiological concentrations, that is it mimics the ion composition of the extracellular fluid. The solution has an osmotic pressure varying from 280 to 340 mOsmol, and a pH varying from 7.0 to 7.4. The buffered saline solution can be selected from the group consisting of Krebs-Henseleit's, Krebs-Ringer's or Hank's buffer, as examples.

The composition of the invention could be suitable to treat and/or prevent diseases such as cardiovascular diseases or treat an injury to heart tissues. Cardiovascular diseases which could be treated include cardiac congenital malformations (e.g. cardiac atrophy, cardiac hypertrophy, defective cardiac chamber organization) or dysfunctions that could be caused by

stress conditions during the fetal life or at birth, including ischemic conditions, infections by microorganisms, exposure to teratogenic toxicants, substances or drugs. Cardiovascular diseases which could be treated also include aging-related heart pathologies, such as heart infarction, congestive heart failure, and acute myocardial ischemia.

The composition could also be involved in modulating heart development during embryogenesis by inducing cardiomyogenesis. The composition of the invention may thus be administered during gestation to correct development of the heart.

The composition of the invention may be administered alone or as part of a more complex pharmaceutical composition according the desired use and route of administration. For instance, the composition of the invention could comprise a vector, such as a plasmid or a virus, comprising a DNA sequence coding for native oxytocin, coding for a modified/fusion oxytocin protein having an increased cardiomyogenic activity, or an increased stability. Anyhow, for preparing such compositions, methods well known in the art may be used.

Oxytocin and/or its derivatives may be coupled to a biocompatible polymer (e.g. polyethylene glycol, polyvinyl alcohol) to reduce antigenicity when administered parenterally.

The composition of the invention and/or more complex pharmaceutical compositions comprising the same may be given via various routes of administration. For instance, the composition may be administered in the form of sterile injectable preparations, for example, as sterile injectable aqueous or oleaginous suspensions. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may also be sterile injectable solutions or suspensions in non-toxic parenterally-acceptable diluents or solvents. They may be given parenterally, for example intravenously, intramuscularly or

sub-cutaneously by injection or by infusion. The composition may also be administered per os (e.g. capsules), nasal spray, transdermal delivery (e.g. iontophoresis). Suitable dosages will vary, depending upon factors such as the amount of each of the components in the composition, the desired effect (fast or long term), the disease or disorder to be treated, the route of administration and the age and weight of the individual to be treated.

In an embodiment, the composition of the invention and/or more complex pharmaceutical compositions comprising the same may be given by direct injection into the heart at the site of infarction or injury. Indeed, damaged sites were shown to attract newly added cardiomyocytes or progenitor cells.

Oxytocin or a functional derivative thereof could also be used in methods for culturing cardiac cells *in vitro*. By providing an effective amount of oxytocin to *in vitro* cultured stem/progenitor cells, it will induce the differentiation of the cultured stem/progenitor cells into cardiac cells, such as cardiomyocytes, and then will promote the aggregation of cardiac cells and promote the tissular organization of *in vitro* cultured heart tissues. Oxytocin or a functional derivative thereof could thus be very useful for providing cardiac tissues for transplant purposes.

Therefore, a related aspect of the invention relates to a method for inducing cardiomyogenic differentiation from cells, such as non-cardiomyocytes (e.g. stem cells), i.e. to induce differentiation of a non-cardiomyocyte to a cardiomyocyte. In a preferred embodiment, the method comprises the step of contacting the non-cardiomyocyte with an effective amount of oxytocin or an oxytocin-related compound or functional derivatives thereof. The method may in embodiments comprise introducing into a cell a nucleic acid (e.g. in a suitable vector) capable of encoding oxytocin or an oxytocin-related compound or functional derivative thereof. According

to another aspect, the present invention provides a method to stimulate the fusion of newly-differentiated cardiomyocytes. The cells are contacted with about  $10^{-10}$  M to about  $10^{-4}$  M of OT, preferably from about  $10^{-9}$  M to about  $10^{-6}$  M of OT, in  
5 embodiments for about 8 h to about 14 days.

The invention further provides a culture medium to induce differentiation of non-cardiomyocytes cultured therein to cardiomyocytes. The culture medium comprises oxytocin or an oxytocin-related compound, or a functional derivative thereof.

10 The oxytocin may in embodiments be present in the medium at a concentration equal to or greater than about  $10^{-10}$  M, in a further embodiment of about  $10^{-10}$  M to about  $10^{-4}$  M, in a further embodiment from about  $10^{-9}$  M to about  $10^{-6}$  M, in yet a further embodiment from about  $10^{-8}$  M to about  $10^{-6}$  M (in  
15 embodiments, about  $10^{-7}$  M of oxytocin for *in vivo* applications and about  $10^{-6}$  M of oxytocin for *ex vivo* applications). The medium may further comprise various standard media components and elements conducive to cell culture, such as salts, acids and bases to control ionic strength and pH of the  
20 medium, antibiotics to reduce contamination, and any other elements or factors conducive to cell culture. Such standard media components are known in the art and are commercially available. The medium may be provided in liquid or powder form, and is in an embodiment substantially sterile. The  
25 medium may be provided in a commercial package together with instructions for differentiation of a non-cardiomyocyte to a cardiomyocyte. The invention further provides a commercial package comprising a standard or base medium, oxytocin or an oxytocin-related compound, or functional derivatives thereof,  
30 together with instructions for differentiation of a non-cardiomyocyte to a cardiomyocyte. The invention further provides a commercial package comprising oxytocin or an oxytocin-related compound, or a functional derivative thereof, together with instructions for differentiation of a non-  
35 cardiomyocyte to a cardiomyocyte.



It will be understood by one skilled in the art that the methods and compositions contemplated by the present invention when applicable, may advantageously be used either *in vitro*, *ex vivo* and/or *in vivo*.

5 With regard to increasing or upregulating expression of a oxytocin in a cell, various methods of introducing oxytocin-encoding nucleic acids into the cell may be used, examples of which are described below. Methods such as the gene therapy methods discussed below may be used in this regard. Examples  
10 of oxytocin-encoding nucleic acids include the nucleic acid of SEQ ID NO:5, a nucleic acid capable of encoding the polypeptide of SEQ ID NOs: 1,2 or 6, or nucleic acids substantially identical thereto. The method may also comprise administering to an area or cardiac tissue a cell comprising such an  
15 oxytocin-encoding nucleic acid, via for example implantation or introduction of such a cell comprising such a oxytocin-encoding nucleic acid.

"Homology" and "homologous" refers to sequence similarity between two peptides or two nucleic acid molecules. Homology  
20 can be determined by comparing each position in the aligned sequences. A degree of homology between nucleic acid or between amino acid sequences is a function of the number of identical or matching nucleotides or amino acids at positions shared by the sequences. As the term is used herein, a nucleic  
25 acid sequence is "homologous" to another sequence if the two sequences are substantially identical and the functional activity of the sequences is conserved (as used herein, the term 'homologous' does not infer evolutionary relatedness).

Two nucleic acid sequences are considered substantially  
30 identical if, when optimally aligned (with gaps permitted), they share at least about 50% sequence similarity or identity, or if the sequences share defined functional motifs. In alternative embodiments, sequence similarity in optimally aligned substantially identical sequences may be at least 60%,  
35 70%, 75%, 80%, 85%, 90% or 95%. As used herein, a given

percentage of homology between sequences denotes the degree of sequence identity in optimally aligned sequences. An "unrelated" or "non-homologous" sequence shares less than 40% identity, though preferably less than about 25 % identity, with a sequence of interest.

Substantially complementary nucleic acids are nucleic acids in which the "complement" of one molecule is substantially identical to the other molecule. Optimal alignment of sequences for comparisons of identity may be conducted using a variety of algorithms, such as the local homology algorithm of Smith and Waterman, 1981, *Adv. Appl. Math* 2: 482, the homology alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, the search for similarity method of Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85: 2444, and the computerised implementations of these algorithms (such as GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, Madison, WI, U.S.A.). Sequence identity may also be determined using the BLAST algorithm, described in Altschul et al., 1990, *J. Mol. Biol.* 215:403-10 (using the published default settings). Software for performing BLAST analysis may be available through the National Center for Biotechnology Information (through the internet at <http://www.ncbi.nlm.nih.gov/>). The BLAST algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold. Initial neighbourhood word hits act as seeds for initiating searches to find longer HSPs. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction is halted when the following

parameters are met: the cumulative alignment score falls off by the quantity  $X$  from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters  $W$ ,  $T$  and  $X$  determine the sensitivity and speed of the alignment. The BLAST program may use as defaults a word length ( $W$ ) of 11, the BLOSUM62 scoring matrix (Henikoff and Henikoff, 1992, *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments ( $B$ ) of 50, expectation ( $E$ ) of 10 (or 1 or 0.1 or 0.01 or 0.001 or 0.0001),  $M=5$ ,  $N=4$ , and a comparison of both strands. One measure of the statistical similarity between two sequences using the BLAST algorithm is the smallest sum probability ( $P(N)$ ), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. In alternative embodiments of the invention, nucleotide or amino acid sequences are considered substantially identical if the smallest sum probability in a comparison of the test sequences is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

An alternative indication that two nucleic acid sequences are substantially complementary is that the two sequences hybridize to each other under moderately stringent, or preferably stringent, conditions. Hybridization to filter-bound sequences under moderately stringent conditions may, for example, be performed in 0.5 M  $\text{NaHPO}_4$ , 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.2 x SSC/0.1% SDS at 42°C (see Ausubel, et al. (eds), 1989, *Current Protocols in Molecular Biology*, Vol. 1, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., New York, at p. 2.10.3). Alternatively, hybridization to filter-bound sequences under stringent conditions may, for example, be performed in 0.5 M  $\text{NaHPO}_4$ , 7% SDS, 1 mM EDTA at 65°C, and washing in 0.1 x SSC/0.1% SDS at 68°C (see Ausubel, et al. (eds), 1989, *supra*).

Hybridization conditions may be modified in accordance with known methods depending on the sequence of interest (see Tijssen, 1993, *Laboratory Techniques in Biochemistry and Molecular Biology -- Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, New York). Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point for the specific sequence at a defined ionic strength and pH.

10       The invention further provides a composition for the prevention and/or treatment of a disease characterized by cardiomyocyte loss or deficiency comprising oxytocin or an oxytocin-related compound, in admixture with a pharmaceutically acceptable carrier.

15       The invention further provides a use of oxytocin or an oxytocin-related compound, or the above-mentioned composition, for the prevention and/or treatment of a disease characterized by cardiomyocyte loss or deficiency.

20       The invention further provides a use of oxytocin or an oxytocin-related compound for the preparation of a medicament for the prevention and/or treatment of a disease characterized by cardiomyocyte loss or deficiency.

25       The invention further provides commercial packages comprising an oxytocin, or an oxytocin-related compound, or the above-mentioned composition together with instructions for the prevention and/or treatment of a disease characterized by cardiomyocyte loss or deficiency.

30       In accordance with another aspect of the invention, therapeutic compositions of the present invention, comprising oxytocin, or an oxytocin-related compound, may be provided in containers or commercial packages which further comprise instructions for their use for the prevention and/or treatment of a disease characterized by cardiomyocyte loss or deficiency.

35       Given that induction of oxytocin receptor (OTR) activity results in an induction of differentiation of a non-

cardiomyocyte to a cardiomyocyte as shown herein, compounds which are capable of modulating OTR expression or activity can be used to induce differentiation of a non-cardiomyocyte to a cardiomyocyte and/or for the prevention and treatment of disease characterized by cardiomyocyte loss or deficiency.

Therefore, the invention further relates to screening methods for the identification and characterization of compounds useful for the induction of differentiation of a non-cardiomyocyte to a cardiomyocyte and/or for the prevention and treatment of disease characterized by cardiomyocyte loss or deficiency.

Therefore, the invention further provides a method of identifying or characterizing a compound for inducing differentiation of a non-cardiomyocyte cell into a cardiomyocyte, or for treatment of a disease characterized by cardiomyocyte loss or deficiency, said method comprising:

(a) contacting a test compound with a cell comprising an oxytocin receptor (OTR) or an OTR-encoding nucleic acid; and

(b) determining whether OTR activity or expression is increased in the presence of the test compound, said increase in OTR activity or expression being an indication that the test compound may be used for inducing differentiation of a non-cardiomyocyte cell into a cardiomyocyte, or for treatment of a disease characterized by cardiomyocyte loss or deficiency.

In an embodiment, the above-mentioned cell comprising OTR is a cell which comprises endogenous levels or expression of OTR. The OTR-comprising cell may be an appropriate host cell in which an exogenous source of OTR was introduced. Such a host cell may be prepared by the introduction of nucleic acid sequences encoding OTR into the host cell under providing conditions for the expression of an OTR. In an embodiment, such a nucleic acid is DNA. Such host cells may be prokaryotic or eukaryotic, bacterial, yeast, amphibian or mammalian. In an embodiment, such host cells are human. DNA and polypeptide

sequences corresponding to human OTR are for example set forth in Figure 12 and SEQ ID NOs: 7 and 8.

The invention further provides a method of identifying or characterizing cells or progenitor cells capable of or having the potential to differentiate into cardiomyocytes, the method comprising determining whether such a cell expresses OTR and/or responds to OT by increasing OTR expression, wherein OTR expression and/or such a response to OT indicate that the cell is capable of or has the potential to differentiate into a cardiomyocyte. In embodiments, the latter method may be used in conjunction with a differentiation method of the invention, whereby a cell is first identified as being capable of differentiation, and is subsequently differentiated into a cardiomyocyte using the methods and agents described herein.

OTR expression may be measured on the transcriptional or translational level, e.g. by the amount of RNA or protein produced. OTR expression can thus be evaluated by RT-PCR, immunocytochemistry, Western blotting (Paquin et al., PNAS 2002) and/or binding studies. RNA may be detected by for example Northern analysis or by the reverse transcriptase-polymerase chain reaction (RT-PCR) method (see for example Sambrook et al (1989) *Molecular Cloning: A Laboratory Manual* (second edition), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA). Protein levels may be detected either directly using affinity reagents (e.g. an antibody or fragment thereof [for methods, see for example Harlow, E. and Lane, D (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY]; a ligand which binds the protein), or by measurement of the protein's activity or a related detectable phenotype.

The above-mentioned method may be employed either with a single test compound or a plurality or library (e.g. a combinatorial library) of test compounds. In the latter case, synergistic effects provided by combinations of compounds may also be identified and characterized. The above-mentioned

compounds may be used for prevention and/or treatment of a disease characterized by cardiomyocyte loss or deficiency, or may be used as lead compounds for the development and testing of additional compounds having improved specificity, efficacy and/or pharmacological (e.g. pharmacokinetic) properties. In certain embodiments, one or a plurality of the steps of the screening/testing methods of the invention may be automated.

In another aspect of the invention, the isolated nucleic acid, for example a nucleic acid sequence encoding oxytocin or a homolog, fragment or variant thereof, may further be incorporated into a recombinant expression vector. In an embodiment, the vector will comprise transcriptional regulatory element or a promoter operably-linked to an oxytocin-coding region. A first nucleic acid sequence is "operably-linked" with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably-linked to a coding sequence if the promoter affects the transcription or expression of the coding sequences.

Generally, operably-linked DNA sequences are contiguous and, where necessary to join two protein coding regions, in reading frame. However, since for example enhancers generally function when separated from the promoters by several kilobases and intronic sequences may be of variable lengths, some polynucleotide elements may be operably-linked but not contiguous. "Transcriptional regulatory element" is a generic term that refers to DNA sequences, such as initiation and termination signals, enhancers, and promoters, splicing signals, polyadenylation signals which induce or control transcription of protein coding sequences with which they are operably-linked. The recombinant expression may further encode additional sequences, such as signal peptide sequences to allow entry of the encoded polypeptide into the secretory pathway, and other domains, which may provide added stability to the polypeptide or are conducive to isolation/purification of the

peptide produced (e.g. fusions with commonly used domains/peptides/affinity tags).

The recombinant expression vector of the present invention can be constructed by standard techniques known to one of ordinary skill in the art and found, for example, in Sambrook et al. (1989) in Molecular Cloning: A Laboratory Manual. A variety of strategies are available for ligating fragments of DNA, the choice of which depends on the nature of the termini of the DNA fragments and can be readily determined by persons skilled in the art. The vectors of the present invention may also contain other sequence elements to facilitate vector propagation and selection in bacteria and host cells. In addition, the vectors of the present invention may comprise a sequence of nucleotides for one or more restriction endonuclease sites. Coding sequences such as for selectable markers and reporter genes are well known to persons skilled in the art. An example of suitable vector is the HSV amplicon-based vector described in the Examples below.

A recombinant expression vector comprising a nucleic acid sequence of the present invention may be introduced into a host cell, which may include a living cell capable of expressing the protein coding region from the defined recombinant expression vector. The living cell may include both a cultured cell and a cell within a living organism. Accordingly, the invention also provides host cells containing the recombinant expression vectors of the invention. The terms "host cell" and "recombinant host cell" are used interchangeably herein. Such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

Vector DNA can be introduced into cells via conventional transformation or transfection techniques. The terms



"transformation" and "transfection" refer to techniques for introducing foreign nucleic acid into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, electroporation, microinjection and viral-mediated transfection. Suitable methods for transforming or transfecting host cells can for example be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory manuals. Methods for introducing DNA into mammalian cells *in vivo* are also known, and may be used to deliver the vector DNA of the invention to a subject for gene therapy for a disease characterized by cardiomyocyte loss or deficiency.

A cell, tissue, organ, or organism into which has been introduced a foreign nucleic acid, is considered "transformed", "transfected", or "transgenic". A transgenic or transformed cell or organism also includes progeny of the cell or organism and progeny produced from a breeding program employing a transgenic organism as a parent and exhibiting an altered phenotype resulting from the presence of a recombinant nucleic acid construct. A transgenic organism is therefore an organism that has been transformed with a heterologous nucleic acid, or the progeny of such an organism that includes the transgene.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (such as resistance to antibiotics) may be introduced into the host cells along with the gene of interest. Preferred selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acids encoding a selectable marker may be introduced into a host cell on the same vector as that encoding the peptide compound or may be introduced on a separate vector. Cells stably transfected

with the introduced nucleic acid may be identified by drug selection (cells that have incorporated the selectable marker gene will survive, while the other cells die).

Given that administration of oxytocin induces  
5 differentiation of a non-cardiomyocyte to a cardiomyocyte, as described herein, a further aspect of the present invention is the treatment of a disease characterized by cardiomyocyte loss or deficiency by administering to a subject a nucleic acid molecule encoding oxytocin or an oxytocin-related compound.  
10 Suitable methods of administration include gene therapy methods.

A nucleic acid of the invention may be delivered to cells in vivo using methods such as direct injection of DNA, receptor-mediated DNA uptake, viral-mediated transfection or  
15 non-viral transfection and lipid based transfection, all of which may involve the use of gene therapy vectors. Direct injection has been used to introduce naked DNA into cells in vivo (see e.g., Acsadi et al. (1991) Nature 332:815-818; Wolff et al. (1990) Science 247:1465-1468). A delivery apparatus  
20 (e.g., a "gene gun") for injecting DNA into cells in vivo may be used. Such an apparatus may be commercially available (e.g., from BioRad). Naked DNA may also be introduced into cells by complexing the DNA to a cation, such as polylysine, which is coupled to a ligand for a cell-surface receptor (see for  
25 example Wu, G. and Wu, C. H. (1988) J. Biol. Chem. 263:14621; Wilson et al. (1992) J. Biol. Chem. 267:963-967; and U.S. Pat. No. 5,166,320). Binding of the DNA-ligand complex to the receptor may facilitate uptake of the DNA by receptor-mediated endocytosis. A DNA-ligand complex linked to adenovirus capsids  
30 which disrupt endosomes, thereby releasing material into the cytoplasm, may be used to avoid degradation of the complex by intracellular lysosomes (see for example Curiel et al. (1991) Proc. Natl. Acad. Sci. USA 88:8850; Cristiano et al. (1993) Proc. Natl. Acad. Sci. USA 90:2122-2126).

35 Defective retroviruses are well characterized for use as

gene therapy vectors (for a review see Miller, A. D. (1990) Blood 76:271). Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology, Ausubel, F. M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines include .psi.Crip, .psi.Cre, .psi.2 and .psi.Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells, endothelial cells, lymphocytes, myoblasts, hepatocytes, bone marrow cells, *in vitro* and/or *in vivo* (see for example Eglitis, et al. (1985) Science 230:1395-1398; Danos and Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:6460-6464; Wilson et al. (1988) Proc. Natl. Acad. Sci. USA 85:3014-3018; Armentano et al. (1990) Proc. Natl. Acad. Sci. USA 87:6141-6145; Huber et al. (1991) Proc. Natl. Acad. Sci. USA 88:8039-8043; Ferry et al. (1991) Proc. Natl. Acad. Sci. USA 88:8377-8381; Chowdhury et al. (1991) Science 254:1802-1805; van Beusechem et al. (1992) Proc. Natl. Acad. Sci. USA 89:7640-7644; Kay et al. (1992) Human Gene Therapy 3:641-647; Dai et al. (1992) Proc. Natl. Acad. Sci. USA 89:10892-10895; Hwu et al. (1993) J. Immunol. 150:4104-4115; U.S. Pat. No. 4,868,116; U.S. Pat. No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573).

For use as a gene therapy vector, the genome of an adenovirus may be manipulated so that it encodes and expresses a peptide compound of the invention, but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. See for example Berkner et al. (1988) BioTechniques 6:616; Rosenfeld et al. (1991) Science 252:431-434; and Rosenfeld et al. (1992) Cell 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or

other strains of adenovirus (e.g., Ad2, Ad3, Ad7 etc.) are well known to those skilled in the art. Recombinant adenoviruses are advantageous in that they do not require dividing cells to be effective gene delivery vehicles and can be used to infect a wide variety of cell types, including airway epithelium (Rosenfeld et al. (1992) cited supra), endothelial cells (Lemarchand et al. (1992) Proc. Natl. Acad. Sci. USA 89:6482-6486), hepatocytes (Herz and Gerard (1993) Proc. Natl. Acad. Sci. USA 90:2812-2816) and muscle cells (Quantin et al. (1992) Proc. Natl. Acad. Sci. USA 89:2581-2584).

Adeno-associated virus (AAV) may be used as a gene therapy vector for delivery of DNA for gene therapy purposes. AAV is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle (Muzyczka et al. Curr. Topics in Micro. and Immunol. (1992) 158:97-129). AAV may be used to integrate DNA into non-dividing cells (see for example Flotte et al. (1992) Am. J. Respir. Cell. Mol. Biol. 7:349-356; Samulski et al. (1989) J. Virol. 63:3822-3828; and McLaughlin et al. (1989) J. Virol. 62:1963-1973). An AAV vector such as that described in Tratschin et al. (1985) Mol. Cell. Biol. 5:3251-3260 may be used to introduce DNA into cells (see for example Hermonat et al. (1984) Proc. Natl. Acad. Sci. USA 81:6466-6470; Tratschin et al. (1985) Mol. Cell. Biol. 4:2072-2081; Wondisford et al. (1988) Mol. Endocrinol. 2:32-39; Tratschin et al. (1984) J. Virol. 51:611-619; and Flotte et al. (1993) J. Biol. Chem. 268:3781-3790). Lentiviral gene therapy vectors may also be adapted for use in the invention.

General methods for gene therapy are known in the art. See for example, U.S. Pat. No. 5,399,346 by Anderson et al.

A biocompatible capsule for delivering genetic material is described in PCT Publication WO 95/05452 by Baetge et al. Methods of gene transfer into hematopoietic cells have also previously been reported (see Clapp, D. W., et al., Blood 78:

1132-1139 (1991); Anderson, Science 288:627-9 (2000); and , Cavazzana-Calvo et al., Science 288:669-72 (2000)).

The invention further relates to implantation, transplantation and grafting methods, to introduce into a  
5 subject a cell comprising a nucleic acid capable of encoding oxytocin or an oxytocin-related compound. The nucleic acid may be present in a vector as described above, the vector being introduced into the cell *in vitro*, using for example the methods described above. In an embodiment, the cell is  
10 autologous, and is obtained from the subject. In embodiments, the cell is non-autologous, e.g. allogenic or xenogenic.

#### EXAMPLES

15 Oxytocin (OT), a nonapeptide largely expressed in the hypothalamus, has long been recognized as a female reproductive hormone necessary for uterine contraction during parturition, timing and amplification of labour, milk ejection during lactation, and ovulation (1). However, the last decades have  
20 shed new light on OT functions. It has been shown that both sexes have equivalent concentrations of OT in the hypophysis and plasma as well as a similar number of oxytocinergic neurons in the hypothalamus (2), and respond to the same stimuli for OT release (3, 4). It also appears that reproductive functions  
25 and maternal behaviour are preserved in OT<sup>-/-</sup> mutant mice (5). Both OT<sup>-/-</sup> males and females are fertile, and females are capable of parturition although they lack the milk ejection reflex (5, 6). These observations indicate that OT is not essential for reproduction, and data now underline the  
30 involvement of OT in sexual behaviour, cognition, memory, tolerance, adaptation, food and water intake, and cardiovascular functions (1, 7, 8).

Recently, a new role has been suggested for OT as a growth and cellular differentiation factor. The

antiproliferative effect of OT, mediated by OT receptors (OTR), has been documented in breast cancer cells (9) and other tumors (10-12). In contrast to its effect on tumoral cells, a mitogenic action of OT has also been described. OT stimulates the proliferation of thymocytes (13, 14) and mitotic activity in the prostate epithelium (15), vascular endothelium (16) and trophoblasts (17). OT has also been reported to enhance myoepithelial cell differentiation and proliferation in the mouse mammary gland (18). The possibility that OT has trophic effects on the embryo has not been investigated intensively. However, OT has been shown to have an influence on the developing heart: OT administered in excess to the fetus may impair cardiac growth in humans and rats (19, 20), and OTR suppression by specific OT antagonists (OTA) in the early stage of chicken egg development leads to cardiac malformation in the embryos (21). It is not known whether the trophic effects of OT on the heart are direct or indirect.

OT's indirect actions could be related to its cardiovascular functions observed in adult rats (7, 22-24). Indeed, we uncovered the entire OT/OTR system in the rat heart, and showed that cardiac OTR activation is coupled to the release of atrial natriuretic peptide (ANP), a potent diuretic, natriuretic and vasorelaxant hormone that is also involved in cell growth regulation (7, 8). A role for ANP in cardiomyogenesis has even been suggested by Cameron et al. (25). In support of a potential action of OT on cardiac development, a maximal OT protein level was seen in the heart at day 21 of gestation and postnatal days 1-4, when cardiac myocytes are at a stage of intense hyperplasia (26).

The P19 mouse embryonal carcinoma cell line is an established model of cell differentiation. Developmentally, pluripotent P19 cells give rise to the formation of cell derivatives of all 3 germ layers (27) (28) and appear to differentiate via the same mechanisms as normal embryonic stem cells (27, 29). When cultured in the presence of  $10^{-6}$  M

retinoic acid (RA), a physiologically-relevant morphogen, P19 cells efficiently ( $\geq 95\%$ ) differentiate to neurons (27, 30, 31). The solvent DMSO induces cardiac differentiation, albeit not as efficiently ( $\leq 15\%$ ) (27, 32). DMSO has been shown to  
5 activate essential cardiogenic transcription factors, such as GATA-4 and Nkx-2.5 (32, 33). However, the mechanisms responsible for triggering these genes in the embryo are still unknown, as is the mode of action of DMSO with respect to the cardiomyogenic program in P19 cells.

10 In the present example, the inventors investigated whether OT induces differentiation of P19 cells into a cardiomyocyte phenotype. The results confirm that OT has a potential naturally-occurring cardiomorphogen activity.

#### 15 **Example 1: Materials And Methods**

##### *Culture and differentiation of P19 cells*

P19 cells were propagated and differentiated according to the procedures of Rudnicki and McBurney (28), with minor  
20 modifications. Undifferentiated cells were propagated in complete medium containing  $\alpha$ -modified Eagle's minimal essential medium ( $\alpha$ -MEM; GIBCO-BRL Burlington, Ontario, Canada) supplemented with 2.5% heat-inactivated fetal bovine serum, 7.5% heat-inactivated donor bovine serum (Cansera  
25 International, Rexdale, Ontario, Canada), and the antibiotics (GIBCO-BRL) penicillin G (50 U/ml) and streptomycin (50  $\mu$ g/ml). The cultures were maintained at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> and passaged every 2 days. The general protocol used for differentiation of P19 cells is depicted in Figure 1.

30 Differentiation was routinely induced with DMSO. Briefly,  $0.25 \times 10^6$  cells were allowed to aggregate for 4 days in non-adhesive bacteriological grade petri dishes (6-cm diameter) containing 5 ml complete medium, in the presence of 0.5% (v/v) DMSO (Sigma Chemical Co., St. Louis, MO). At day 2 of  
35 aggregation, the inducing culture medium was replenished. At

day 4, aggregates were transferred to tissue culture grade vessels (10-cm diameter dishes or 24/48-well plates), and cultured in complete medium in the absence of differentiation-inducing agent. Aggregation was also done in the absence of DMSO, and in the presence of  $10^{-7}$  M OT and/or  $10^{-7}$  OTA ([d(CH<sub>2</sub>)<sub>5</sub><sup>1</sup>, Tyr(Me)<sup>2</sup>, Thr<sup>4</sup>, Orn<sup>8</sup>, Tyr-NH<sub>2</sub><sup>9</sup>]-vasotocin), both from Peninsula Laboratories Inc. (San Carlos, CA). The cell populations were analyzed at days 10-14 of the entire differentiation protocol, at a time cardiac cells normally beat synchronously.

#### *Cell morphology, staining and immunocytochemistry*

Examinations were done under a Zeiss® inverted microscope (Zeiss IM, Carl Zeiss, Jena, Germany) equipped with phase-contrast objectives, filters for rhodamine and fluorescein fluorescence, a MC 100® camera and a photoautomat unit. Micrographs were taken with Kodak Technical Pan® film (for cell morphology) or with Kodak T-Max 400® or Elite-II 100® film (for fluorescence).

For morphological examination, cells were grown directly onto the plastic surface of tissue culture vessels. For staining with rhodamine<sup>123</sup> (Sigma), day-4 aggregates were distributed in 24-well culture plates and grown until day 8. Then, dye was added to the culture medium at a final concentration of 1 µg/ml for 45 min, and afterwards, the cells were washed extensively with phosphate-buffered saline (PBS) and cultured for 48 h in the absence of the dye. Dye retained by cells in each well was measured by a fluorescence microplate reader (SPECTRA Max Gemini®, Molecular Devices, Sunnyvale, CA) at 505 nm for excitation and 534 nm for emission.

For immunocytofluorescence studies, cells were grown onto glass coverslips coated with 0.1% gelatin. They were then fixed by 20-min incubation in PBS containing 4% paraformaldehyde, rinsed in PBS and stored at 4°C in this buffer until used. All subsequent steps of permeabilization, washing and incubation



with antibodies were performed at room temperature. Fixed cells were permeabilized for 10 min in PBS containing 0.005% saponin, blocked for 60 min in PBS-BSA-saponin (PBS containing 1% bovine serum albumin and 0.005% saponin), incubated for 45 min with the primary antibody diluted 1/50 and for 45 min with a fluorescein-conjugated swine anti-goat IgG antibody (Biosource International, Camarillo, CA) diluted 1/1000. PBS-BSA-saponin was used for washing between incubations and antibodies were diluted in the same buffer but containing 1.5% normal swine serum (Jackson Immuno Research Laboratories Inc., West Grove, PA). Coverslips were mounted in PBS containing 50% glycerol, and immediately examined under the microscope. The primary antibodies were all from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA) and produced in goat: antibody C-20 against OT receptor (OTR), antibody K-16 against sarcomeric myosin heavy chain (MHC), and antibody N-19 against dihydropyridine receptor-alpha1 (DHPR-alpha1).

#### *Analysis of OT by radioimmunoassay (RIA)*

P19 cells induced with DMSO or OT were cultured until day 14 in absence of inducer. Cells were washed carefully with PBS and incubated for 4 h in serum-free  $\alpha$ -MEM supplemented with protease inhibitors (aprotinin, 30  $\mu$ g/mL and soybean trypsin inhibitor (SBTI), 100  $\mu$ g/mL). Cells and media were collected, extracted on Sep-Pak C<sub>18</sub>-cartridges (Waters Canada, Mississauga, Ontario), lyophilized, resuspended in a minimal volume of  $\alpha$ -MEM and assayed by RIA as described (8). RIA was also done on undiluted serum combination used for P19 cell culture (1 part of fetal bovine serum and 3 parts of donor bovine serum).

#### *Analysis by reverse transcription-polymerase chain reaction (RT-PCR)*

Total cellular RNA was extracted with TRIzol® Reagent (Invitrogen Life Technologies, Burlington, Ontario, Canada),

and poly(A)<sup>+</sup>mRNA was affinity purified from 200 µg of total RNA onto Oligotex® mRNA columns (Qiagen, Mississauga, Ontario, Canada), as per the manufacturers' instructions. First-strand cDNA was synthesized in a final volume of 40 µl containing

5 first-strand buffer, 3 µg of cellular RNA, 4 µl of hexanucleotide primers (Amersham-Pharmacia, Baie d'Urfé, Quebec, Canada), and avian myeloblastosis virus reverse transcriptase (12 units/µg RNA; Invitrogen). First-strand cDNA (5 µl) was then used for PCR amplification with OTR, ANP or

10 GAPDH exon-specific oligonucleotide primers in a Robocycler Gradient 40 thermocycler (Stratagene, La Jolla, CA). Sequences of mouse OTR and ANP genes have been described (26, 34). Conditions for RT-PCR analysis of mouse OTR were adapted from Wagner et al. (6, 7). For all PCR studies the number of cycles

15 used was within the linear range of amplification. The OTR sense and antisense primers were respectively the 22-bp 5'-AAGATGACCTTCATCATTTGTTTC-3' (SEQ ID NO:9) and the 23-bp 5'-CGACTCAGGACGAAGGTGGAGGA-3' (SEQ ID NO:10). Amplification was performed over 32 cycles, each involving 1 min at 94°C, 1.5 min

20 at 62°C and 1.5 min at 72°C, and was terminated by a 5-min final extension at 72°C. The ANP antisense and sense primers were respectively the 24-bp 5'-GTCAATCCTACCCCGAAGCAGCT-3' (SEQ ID NO:11) and the 20-bp 5'-CAGCATGGGCTCCTTCTCCA-3' (SEQ ID NO:12). Amplification was performed over 25-30 cycles, each

25 involving 1 min at 94°C, 1 min at 65°C and 3 min at 72°C, and was terminated by a 5-min final extension at 72°C. The amplification of GAPDH mRNA, a constitutively and ubiquitously expressed gene, served as an internal standard for RT-PCR analysis.

30 The 23-bp antisense primer 5'-CAGTGATGGCATCCACTGTGGTC-3' (SEQ ID NO:13) and the 23-bp sense primer 5'-AAGGTCGGTGTCAACCCATTTGGCCGT-3' (SEQ ID NO:14) were used. Amplification was performed over 23 cycles, each involving 1 min at 94°C, 1.5 min at 59°C and 2 min at 72°C.

### *Western blot analysis*

Cells were collected by scraping, homogenized in sucrose buffer (20 mM Hepes/Tris, pH 7.4, containing 250 mM sucrose and 20 µg/ml of the protease inhibitor phenylmethylsulfonyl fluoride), then centrifuged at 3000 g for 10 min at 4°C to remove debris. The supernatants were centrifuged at 100 000 g for 45 min at 4°C, and the pellets were resuspended in sucrose buffer for analysis of protein content by a modified Bradford assay (30). Aliquots (20 µg protein) were subjected to polyacrylamide gel electrophoresis in the presence of sodium dodecyl sulfate (SDS-PAGE) under reducing conditions (35) followed by electrotransfer onto pure nitrocellulose membrane (Hybond-C; Amersham-Pharmacia). Molecular size calibration was achieved using Broad Standard Solution (Bio-Rad Laboratories Ltd., Mississauga, Ontario, Canada). The nitrocellulose blots were blocked overnight with 5% nonfat milk in Tris-buffered saline (TBS: 20 mM Tris·Cl, pH 8.0, 140 mM NaCl, 1% BSA and 0.1% Tween-20), then probed with goat C20 antibody (anti-OTR; 1/1,000) for 2 h at room temperature. Antibody incubations and washes were performed in TBS throughout. Detection was realized by enhanced chemiluminescence with an Amersham-Pharmacia ECL® kit and an appropriate peroxidase-conjugated secondary antibody (29). Autoluminograms were developed in an AFP Imaging Mini-med 190® X-Ray Film Processor (AFP Corp., Elmsford, NY).

### *Statistics*

Results are reported as the mean values ± SEM. Comparisons between treatments were done by unpaired Student's t test.

### **Example 2: Effect of OT or DMSO treatment on P19 cells**

Using the time schedule depicted in Figure 1, treatment of P19 cell aggregates with  $10^{-7}$  M OT induced the formation of

rhythmically-beating cells resembling primary cardiomyocytes isolated from the heart of newborn animals. A similar phenotypic change was already reported for treatment with 0.5-1% DMSO (27, 28, 30, 32). Aggregates treated with OT or DMSO were observed to have a 1.5-fold smaller mean diameter than their untreated counterparts (data not shown), a finding that could reflect the antimitotic activity of OT and DMSO.

We examined whether treatment of cell aggregates with OT induced the expression of the cardiac muscle markers sarcomeric MHC and DHPR-alpha1. Sarcomeric MHC is expressed in contractile muscular cells as is DHPR-alpha1, a component of intracellular junctions critical for the coupling of excitation-contraction (27, 32, 36). As presented in Figure 2B, undifferentiated cells were negative for MHC, as reported (27, 28, 32), and for DHPR-alpha1. However as with DMSO, OT induced the appearance of numerous, intense, immunoreactive foci in cell populations (Figs 2A, B). In both cases, there were cell subpopulations that did not respond positively (Fig. 2A) and seemed to be mainly undifferentiated cells according to morphological criteria. We and others have shown that undifferentiated cells remain in DMSO-treated P19 cultures by probing for Stage-specific Embryonic Antigen-1, an established marker of the undifferentiated state (27, 28, 30). Cell aggregates not exposed to OT or DMSO were not positive for MHC and DHPR-alpha1 although they sometimes showed very rare and small immunoreactive foci (Fig. 2B, No inducer). This occasional staining could be due to spontaneous differentiation events triggered by high cell densities such as those encountered in aggregates (27, 28).

We also compared the cardiogenic potency of OT and DMSO. First, potency was simply quantitated by rhodamine<sup>123</sup> retention in cells, taking advantage of the fact that this dye, which penetrates all cell types, is retained for much longer periods (days instead of hours) in cardiac cells than in other cell types (37). To meet their energy requirements for muscular

contraction, cardiomyocytes have indeed abundant mitochondria, the cell organelles that accumulate rhodamine<sup>123</sup>. Figure 3A shows that exposure of the cell aggregates to OT and DMSO significantly increased cellular retention of the dye by 2-3 fold compared to non-induced aggregates ( $p < 0.001$ ), and this increase at day 10 of differentiation was even significantly higher after OT than DMSO treatment ( $p < 0.001$ ). Since P19-derived cardiomyocytes beat in culture, we also compared the time course of appearance of beating cells after treatment of aggregates with DMSO or OT. We found that OT stimulated the production of beating cell colonies in all 24 independently growing cultures by day 8 whereas the same result was obtained in cells induced by DMSO only by day 12 (Fig. 3B). The cardiogenic action of OT was specific and receptor-mediated, since no beating cells were seen when  $10^{-7}$  M OTA was used in place of OT or in combination with OT (Fig. 3B). Interestingly, OTA also abolished the cardiogenic action of DMSO (Fig. 3B). Analysis of OT production by RIA also confirmed cardiomyocyte functioning. Like normal cardiomyocytes, P19 cells induced with DMSO and OT release OT in their culture medium (Fig. 6). Finally, cardiogenic potency was evaluated via ANP expression since this peptide is abundantly produced by cardiomyocytes. The results showed that at day 14 of differentiation ANP mRNA level was significantly upregulated in OT-treated P19 aggregates as compared to undifferentiated cells ( $p < 0.05$ ), and this upregulation was at similar level after DMSO treatment (Fig. 3C). As for cell beating, OTA prevented OT-induced upregulation of ANP expression (Fig. 3C,  $p < 0.05$ ). Although the effect of OTA on DMSO-induced ANP expression was not statistically significant, the inhibitory tendency was observed in all experiments (Fig. 3C). The inhibitory action of OTA on DMSO cardiomyogenic properties was thus more evident by the beating than the ANP criteria. Altogether, rhodamine<sup>123</sup> absorption, and the time-course formation of beating cells and abundance of ANP mRNA pointed to a potent cardiomyogenic effect

of OT. In addition, the cardiomyogenic action of OT and even that of DMSO appear to involve OTR.

To further investigate the involvement of OTR in cardiomyogenesis, we examined OTR expression in P19 cells. OTR protein (Fig. 4A, B) and mRNA (Fig. 4C) were present at low levels in undifferentiated cells, indicating that these cells can respond minimally to OT. OTR expression remained at low levels in aggregates not exposed to OT or DMSO (Fig. 4C, No inducer). In contrast, intense OTR immunoreactive foci were observed in cell populations after OT or DMSO treatment (Fig. 4A). These findings corresponded to the results of Western blotting (Fig. 4B) and RT-PCR analysis of OTR (Fig. 4C), both indicating increased OTR expression. In accordance with the absence of a cardiomyogenic effect of OTA and the inhibitory action of OTA on OT-induced cardiac differentiation, OTA did not upregulate OTR expression by itself and inhibited OT-induced OTR upregulation (Fig. 4B). Thus, the OTR-dependent cardiogenic effect of OT and DMSO seems to involve upregulation of OTR expression.

As described herein, it is shown that OT added to the culture medium of P19 stem cell aggregates induced cardiomyogenic differentiation, which was demonstrated by monitoring the expression of MHC, DHPR- $\alpha$ 1 and ANP cardiac markers, production of OT, retention of a mitochondrial-specific dye and the appearance of beating cell colonies. The cardiogenic effect of OT was specific and mediated by OTR because it was abolished by OTA. OT also upregulated OTR expression. These results suggest a new role for the OT/OTR system in heart genesis and development.

The P19 cell line is an excellent cell differentiation model that mimics the events of early cardioembryogenesis. Differentiation of P19 cells to cardiomyocytes by aggregation and exposure to DMSO was shown to be associated with induction of the cardiac-specific subtype of endothelin receptors (38). In addition, brain natriuretic peptide and ANP were observed

in newly-formed striated muscle structures upon DMSO treatment and not in undifferentiated P19 cells and their neuronal derivatives (39). In the studies described herein, DMSO- and OT-induced ANP transcript levels reached about 5-10% of that found in the adult mouse atrium - the richest site of ANP synthesis. Several transcription factors having an essential role in cardiogenesis are upregulated in DMSO-induced P19 cells. This was shown to be the case for the zinc-finger containing GATA-4, the homeobox gene Nkx2-5, and the myocyte enhancer factor 2C (32, 33, 40), and the overexpression of either factor in P19 cells was sufficient to induce cardiac differentiation in the absence of DMSO (32, 41, 42). Little is known about the molecular mechanisms underlying the activation of these genes, but DMSO was found to increase intracellular  $\text{Ca}^{2+}$  levels and was suspected to affect a pathway that has an extracellular component, possibly serum-borne (27, 43, 44). Interestingly, the data described herein indicate that OTR are upregulated to a similar extent by OT and DMSO, and other studies have reported that OTR function modulates intracellular  $\text{Ca}^{2+}$  concentration in some cell types (1). It is thus tempting to suggest that OT could be a serum-borne factor that is active in DMSO-induced differentiation.

One of the mechanisms by which OT and DMSO trigger cardiac differentiation involves OTR since both agents upregulated the expression of this receptor, and OTA totally abolished their cardiomyogenic action as well as prevented OT-stimulated effect on OTR expression. Homologous regulation of OTR expression by OT itself was observed in the brain and in astroglial cell cultures (46, 47). It is noteworthy that, like DMSO, RA, used at low levels ( $10^{-8}$ - $10^{-9}$  M), induces cardiac differentiation of P19 cells (27, 28). This observation could have some relevance to the OT/OTR system since RA was shown to upregulate OT expression in the fetal heart (26).

Several studies have proposed a role for OT as a growth and differentiation/maturation factor in a

gestational/perinatal context. In the mother, OT is required for postpartum alveolar proliferation, and induces differentiation and proliferation of myoepithelial cells of the mammary gland necessary for milk ejection (1, 18). The OT/OTR system is expressed in human cumulus/luteal cells surrounding oocytes and weak OTR gene expression is even observed in oocytes (48). Moreover, when fertilized mouse oocytes are cultured with OT in vitro, they develop at a higher rate into the blastocyst stage than their unstimulated counterparts (48). Spontaneous myometrial contractures are known to occur during pregnancy in sheep and controlled contractures induced by application of OT pulses to pregnant ewes have been shown to accelerate fetal cardiovascular function (49).

All these studies thus strongly suggest involvement of the maternal and embryonal OT/OTR systems in development of the embryo, and our work points to a particular involvement of OT in the priming of cardiogenesis. We think that OT could also assist the maturation of newly-differentiated cardiomyocytes by stimulating their fusion since beating cells derived from OT-induced P19 cells formed fiber-like structures. Such a fusogenic action was recently reported for OT on skeletal myoblasts in vitro (50). Our results may find application in therapies that consider the replacement of cardiac tissue lost after injury. In this context, OT could be used as a trophic factor to assist the compensatory division of myocytes shown to occur in infarcted organs (51), or to prime the cardiomyogenesis of a variety of progenitor/stem cells to be grafted in the injured heart (52, 53).

### **Example 3: Studies of the mechanism of differentiation mediated by OT/OTR system**

#### **Materials and Methods**



*Induction of P19 cells with different OT concentrations, RA and triiodothyronine ( $T_3$ )*

P19 cells were subjected to the cardiac differentiation protocol depicted in Example 1, except that different concentrations of OT were used during the four days of aggregation. At day 4, aggregates were transferred to 24-well plates and scored for the appearance of beating cell colonies. In addition, cells were collected at day 14 for analysis of OTR or OT expression by RT-PCR. Cell aggregates were also treated with RA and/or  $T_3$ .

*OT RT-PCR*

RT-PCR for OT was performed with three different pairs of primers covering the entire coding sequence of OT gene (8).

OT differentiation of P19 cells into rhythmically beating cardiomyocytes is concentration-dependent within the range of  $10^{-6}$  to  $10^{-9}$  M (Figure 5). OTR mRNA expression in differentiated populations (at day 14) also supports the concentration-dependent effect (Figure 5). Serum was recently recognized as a potent OTR stimulator in various cells (reviewed by Kimura et al. 2003). We have data showing that OT concentration in fetal and donor bovine sera (used in P19 cell cultures) is  $10^{-11}$  M. Consequently, the final OT concentration of  $10^{-12}$  M in the culture medium is not sufficient to induce cardiac differentiation of P19 cells. It is possible that some exogenously supplied compounds, such as DMSO, can induce cardiomyocyte differentiation in conditions of reduced OT concentration (below  $10^{-9}$  M) by increasing OTR affinity for OT, by upregulating OTR expression and/or by inducing the same signaling transducers as OT. Serum-borne factors could also participate in DMSO action.

Retinoic acid (RA), a very well known cardiac morphogen, stimulated P19 cell differentiation into cardiomyocytes at concentration of  $10^{-7}$  M, as reported previously (27,28). This is demonstrated by the appearance of rhythmically beating cells

and OTR upregulation (Figure 7). As in the case of OT and DMSO (Paquin et al., PNAS 2002), OTA completely inhibits RA-induced cardiac differentiation (Figure 7). This new result suggests that OT/OTR system is involved in the cardiomyogenic action of RA. We noticed that the presence of OTA in cultures induced with  $10^{-8}$  M RA (a cardiomyogenic concentration) results in the apparition of some neural cells (Figure 8). The neurogenic potential of RA is known in P19 model but is realized at higher concentration ( $10^{-6}$  M; 27-31). This observation suggests that OT and RA pathways cross-talk during cell differentiation.

The thyroid hormone  $T_3$  was also reported to induce cardiomyogenic differentiation of P19 cells, with an optimal effective concentration of 30 nM (Rodriguez et al., 1994). Similar to the results observed with DMSO and RA-induced differentiation, OTA abolished cardiac differentiation induced by  $T_3$  (Figure 9). Altogether, the results obtained with OTA suggest that OT/OTR could be universally involved in mechanisms of action of cardiomyogenic substances.

P19 stem cells express oxytocin receptor as shown by RT-PCR and immunocytochemistry (Paquin et al., PNAS 2002). Preliminary results suggest that differentiation of stem cells into cardiomyocytes depends on the early potential of inducer to increase OTR. This stimulatory effect is already observed by RT-PCR in the second day of differentiation protocol. In this respect OT is a most potent OTR inducer comparing to RA and DMSO. OT antagonist (OTA) has no effect on OTR expression at this stage of differentiation.

Throughout this application, various references are referred to describe more fully the state of the art to which this invention pertains. The disclosures of these references are hereby incorporated by reference into the present disclosure.

## REFERENCES

1. Gimpl, G. & Fahrenholz, F. (2001) *Physiol Rev.* **81**, 629-683.
2. Ashton, N. & Balment, R. J. (1991) *Acta Endocrinol. (Copenh)* **124**, 91-97.
3. Stoneham, M. D., Everitt, B. J., Hansen, S., Lightman, S. L. & Todd, K. (1985) *J Endocrinol* **107**, 97-106.
4. Verbalis, J. G., Mangione, M. P. & Stricker, E. M. (1991) *Endocrinology* **128**, 1317-1322.
5. Nishimori, K., Young, L. J., Guo, Q., Wang, Z., Insel, T. R. & Matzuk, M. M. (1996) *Proc Natl Acad Sci U S A* **93**, 11699-11704.
6. Wagner, K. U., Young, W. S., III, Liu, X., Ginns, E. I., Li, M., Furth, P. A. & Hennighausen, L. (1997) *Genes Funct.* **1**, 233-244.
7. Gutkowska, J., Jankowski, M., Lambert, C., Mukaddam-Daher, S., Zingg, H. H. & McCann, S. M. (1997) *Proc Natl Acad Sci U S A* **94**, 11704-11709.
8. Jankowski, M., Hajjar, F., Al Kawas, S., Mukaddam-Daher, S., Hoffman, G., McCann, S. M. & Gutkowska, J. (1998) *Proc Natl Acad Sci U S A* **95**, 14558-14563.
9. Cassoni, P., Sapino, A., Fortunati, N., Munaron, L., Chini, B. & Bussolati, G. (1997) *Int. J. Cancer* **72**, 340-344.
10. Cassoni, P., Sapino, A., Stella, A., Fortunati, N. & Bussolati, G. (1998) *Int. J. Cancer* **77**, 695-700.
11. Cassoni, P., Fulcheri, E., Carcangiu, M. L., Stella, A., Deaglio, S. & Bussolati, G. (2000) *J. Pathol.* **190**, 470-477.
12. Copland, J. A., Ives, K. L., Simmons, D. J. & Soloff, M. S. (1999) *Endocrinology* **140**, 4371-4374.
13. Martens, H., Kecha, O., Charlet-Renard, C., Defresne, M. P. & Geenen, V. (1998) *Neuroendocrinology* **67**, 282-289.
14. Geenen, V., Kecha, O., Brilot, F., Charlet-Renard, C. & Martens, H. (1999) *Neuroimmunomodulation*. **6**, 115-125.
15. Plecas, B., Popovic, A., Jovovic, D. & Hristic, M. (1992) *J. Endocrinol. Invest* **15**, 249-253.

16. Thibonnier, M., Conarty, D. M., Preston, J. A., Plesnicher, C. L., Dweik, R. A. & Erzurum, S. C. (1999) *Endocrinology* **140**, 1301-1309.
- 5 17. Cassoni, P., Sapino, A., Munaron, L., Deaglio, S., Chini, B., Graziani, A., Ahmed, A. & Bussolati, G. (2001) *Endocrinology* **142**, 1130-1136.
18. Sapino, A., Macri, L., Tonda, L. & Bussolati, G. (1993) *Endocrinology* **133**, 838-842.
- 10 19. Chard, T., Boyd, N. R., Forsling, M. L., McNeilly, A. S. & Landon, J. (1970) *J Endocrinol* **48**, 223-234.
20. Schrieffer, J. A., Lewis, P. R. & Miller, J. W. (1982) *Biol Reprod* **27**, 362-368.
- 15 21. Widmer, H., Durroux, T., Kempf, H., Mouillac, B., Gasc, J. M. & Barberis, C. (1999) *Abstracts of 1999 World Congress on Neurohypophysial Hormones* 94.
22. Haanwinckel, M. A., Elias, L. K., Favaretto, A. L., Gutkowska, J., McCann, S. M. & Antunes-Rodrigues, J. (1995) *Proc Natl Acad Sci U S A* **92**, 7902-7906.
- 20 23. Favaretto, A. L., Ballejo, G. O., Albuquerque-Araujo, W. I., Gutkowska, J., Antunes-Rodrigues, J. & McCann, S. M. (1997) *Peptides* **18**, 1377-1381.
24. Mukaddam-Daher, S., Lin, Y. L., Gutkowska, J. & Cardinal, R. (2001) *Hypertension*.
- 25 25. Cameron, V. A., Aitken, G. D., Ellmers, L. J., Kennedy, M. A. & Espiner, E. A. (1996) *Endocrinology* **137**, 817-824.
26. Gutkowska, J., Bhat, P., Wang, D., Mukaddam-Daher, S., McCann, S. M. & Jankowski, M. (2002) *19th Scientific meeting of the international society of hypertension, Prague, June 2002 (Abstract)*.
- 30 27. McBurney, M. W. (1993) *Int J Dev Biol* **37**, 135-140.
28. Rudnicki, M. A. & McBurney, M. W. (1987) in *Teratocarcinomas and embryonic stem cells: a practical approach.*, ed. Robertson, E. J. (IRL Press, Oxford, UK), pp. 19-49.
- 35 29. Laplante, I., Paquin, J. & Beliveau, R. (2001) *Brain Res. Dev. Brain Res.* **129**, 157-168.
30. Jeannotte, R., Paquin, J., Petit-Turcotte, C. & Day, R. (1997) *DNA Cell Biol* **16**, 1175-1187.

31. Cadet, N. & Paquin, J. (2000) *Dev Brain Res* **120**, 211-221.
32. Skerjanc, I. S. (1999) *Trends Cardiovasc. Med.* **9**, 139-143.
33. Srivastava, D. & Olson, E. N. (2000) *Nature* **407**, 221-226.
- 5 34. Seidman, C. E., Bloch, K. D., Smith, J. A. & Seidman, J. G. (1984) *Science* **226**, 1206-1209.
35. Laemmli, U. K. (1970) *Nature* **227**, 680-685.
36. Flucher, B. E. & Franzini-Armstrong, C. (1996) *Proc. Natl. Acad. Sci. U. S. A* **93**, 8101-8106.
- 10 37. Summerhayes, I. C., Lampidis, T. J., Bernal, S. D., Nadakavukaren, J. J., Nadakavukaren, K. K., Shepherd, E. L. & Chen, L. B. (1982) *Proc. Natl. Acad. Sci. U. S. A* **79**, 5292-5296.
38. Monge, J. C., Stewart, D. J. & Cernacek, P. (1995) *J Biol Chem* **270**, 15385-15390.
- 15 39. Boer, P. H. (1994) *Biochem. Biophys. Res. Commun.* **199**, 954-961.
40. Mohun, T. & Sparrow, D. (1997) *Curr. Opin. Genet. Dev.* **7**, 628-633.
- 20 41. Grepin, C., Nemer, G. & Nemer, M. (1997) *Development* **124**, 2387-2395.
42. Skerjanc, I. S., Petropoulos, H., Ridgeway, A. G. & Wilton, S. (1998) *J. Biol. Chem.* **273**, 34904-34910.
43. Morley, P. & Whitfield, J. F. (1993) *J. Cell Physiol* **156**, 219-225.
- 25 44. Wilton, S. & Skerjanc, I. (1999) *In Vitro Cell Dev. Biol. Anim* **35**, 175-177.
45. Hartman, R. D., Rosella-Dampman, L. M., Emmert, S. E. & Summy-Long, J. Y. (1986) *Endocrinology* **119**, 1-11.
- 30 46. Insel, T. R., Winslow, J. T. & Witt, D. M. (1992) *Endocrinology* **130**, 2602-2608.
47. Di Scala-Guenot, D. & Strosser, M. T. (1995) *Am. J. Physiol* **268**, C413-C418.
- 35 48. Furuya, K., Mizumoto, Y., Makimura, N., Mitsui, C., Murakami, M., Tokuoka, S., Ishikawa, N., Nagata, I., Kimura, T. & Ivell, R. (1995) *Adv. Exp. Med. Biol.* **395**,

523-528.

49. Shinozuka, N., Yen, A. & Nathanielsz, P. W. (2000) *Am. J. Physiol Heart Circ. Physiol* **278**, H41-H49.
50. Breton, C., Haenggeli, C., Barberis, C., Heitz, F., Bader, C. R., Bernheim, L. & Tribollet, E. (2002) *J. Clin. Endocrinol. Metab* **87**, 1415-1418.
51. Beltrami, A. P., Urbanek, K., Kajstura, J., Yan, S. M., Finato, N., Bussani, R., Nadal-Ginard, B., Silvestri, F., Leri, A., Beltrami, C. A. et al. (2001) *N. Engl. J. Med.* **344**, 1750-1757.
52. Condorelli, G., Borello, U., De Angelis, L., Latronico, M., Sirabella, D., Coletta, M., Galli, R., Balconi, G., Follenzi, A., Frati, G. et al. (2001) *Proc. Natl. Acad. Sci. U. S. A* **98**, 10733-10738.
53. Jackson, K. A., Majka, S. M., Wang, H., Pocius, J., Hartley, C. J., Majesky, M. W., Entman, M. L., Michael, L. H., Hirschi, K. K. & Goodell, M. A. (2001) *J. Clin. Invest* **107**, 1395-1402.
54. Rodriguez ER, Tan CD, Onwuta US, Parrillo JE. 1994. Cardiac myocyte differentiation induced by 3,5,3'-triiodo-L-thyronine (T3) in P19 teratocarcinoma cells is accompanied by preferential binding of RGG(T/A)CA direct repeats spaced by 4 base pairs in the DNA.. *Biochem. Biophys. Res. Commun.* 205:1899-1906).

Although preferred embodiments of the present invention have been described in detail herein and illustrated in the accompanying drawings, it is to be understood that the invention is not limited to these precise embodiments and that various changes and modifications may be effected therein without departing from the scope or spirit of the present invention.


## WHAT IS CLAIMED IS:

1. A method of inducing differentiation of a non-cardiomyocyte into a cardiomyocyte, said method comprising stimulating oxytocin receptor (OTR) activity in said non-cardiomyocyte.

2. The method of claim 1, wherein said method comprises contacting said non-cardiomyocyte with an agent capable of stimulating OTR activity.

3. The method of claim 2, wherein said agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).

4. The method of claim 3, wherein said oxytocin or functional derivative thereof has the structure:



wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.

5. The method of claim 1, wherein the method comprises introducing into the non-cardiomyocyte a nucleic acid capable of encoding oxytocin or an oxytocin-related compound.

6. The method of claim 5, wherein the nucleic acid is selected from the group consisting of:

(a) SEQ ID NO:5;

(b) a nucleic acid sequence capable of

encoding SEQ ID NO:6; and


- (c) a nucleic acid sequence substantially identical to  
(a) or (b).

- 5 7. The method of claim 1, wherein said non-cardiomyocyte is  
a stem or progenitor cell.
8. The method of claim 7, wherein said stem or progenitor  
cell is selected from the group consisting of embryonic  
10 and adult stem or progenitor cells.
9. The method of claim 7, wherein said stem or progenitor  
cell is selected from the group consisting of circulating  
and non-circulating stem or progenitor cells.
- 15 10. The method of claim 7, wherein said method is performed  
*in vitro*.
11. The method of claim 7, wherein said method is performed  
20 *in vivo*.
12. The method of claim 1, wherein said cardiomyocyte is  
characterized by an alteration of a phenotypic feature  
relative to said non-cardiomyocyte, wherein said  
25 phenotypic feature is selected from the group consisting  
of:
- (a) level of oxytocin receptor (OTR) protein or OTR-  
encoding nucleic acid;
- (b) level of ANP protein or ANP-encoding nucleic acid;
- 30 (c) level of muscular MHC protein or muscular MHC-  
encoding nucleic acid;
- (d) level of DHPR-alpha1 protein or DHPR-alpha1-encoding  
nucleic acid;
- (e) level of sarcomeric marker proteins;
- 35 (f) level of ion channels;



- (g) mitochondrial dye retention;
- (h) appearance of rhythmic beats; and
- (i) chronotropic responses.

- 5 13. A method of treating a disease characterized by cardiomyocyte loss or deficiency in an animal, said method comprising stimulating oxytocin receptor (OTR) activity in a non-cardiomyocyte cell of said animal.
- 10 14. The method of claim 13, wherein said method comprises administering an agent capable of stimulating OTR activity to said animal.
- 15 15. The method of claim 14, wherein said agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).
- 20 16. The method of claim 15, wherein said oxytocin or functional derivative thereof has the structure:



- 25 wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.
- 30 17. The method of claim 15, wherein the method comprises administering a nucleic acid capable of encoding oxytocin or a functional derivative thereof to said animal.
18. The method of claim 17, wherein the nucleic acid is selected from the group consisting of:
- (a) SEQ ID NO:5;
  - 35 (b) a nucleic acid sequence capable of

encoding SEQ ID NO:6; and

(c) a nucleic acid sequence substantially identical to (a) or (b).

5 19. The method of claim 13, wherein said non-cardiomyocyte is a stem or progenitor cell.

20. The method of claim 19, wherein said stem or progenitor cell is selected from the group consisting of circulating  
10 and non-circulating stem or progenitor cells.

21. The method of claim 13, wherein said animal is a mammal.

22. The method of claim 13, wherein said animal is a human.  
15

23. The method of claim 13, wherein said disease is selected from the group consisting of cardiac congenital dysfunctions, aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial  
20 ischemia.

24. A method of treating a disease characterized by cardiomyocyte loss or deficiency in an animal, said method comprising:

25 (a) inducing, using the method of claim 1, differentiation of a non-cardiomyocyte cell into a cardiomyocyte; and

(b) implanting said cardiomyocyte into said animal.

30 25. The method of claim 24, wherein said animal is a mammal.

26. The method of claim 24, wherein said animal is a human.


27. The method of claim 24 where said disease is selected from  
35 the group consisting of cardiac congenital dysfunctions,

aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial ischemia.

28. The method of claim 24, wherein said method comprises contacting said non-cardiomyocyte with an agent capable of stimulating OTR activity.

29. The method of claim 28, wherein said agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).

30. The method of claim 29, wherein said oxytocin or functional derivative thereof has the structure:



wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.

31. The method of claim 24, wherein the method comprises introducing into the non-cardiomyocyte a nucleic acid capable of encoding oxytocin or a functional derivative thereof.

32. The method of claim 31, wherein the nucleic acid is selected from the group consisting of:

- (a) SEQ ID NO:5;
- (b) a nucleic acid sequence capable of encoding SEQ ID NO:6; and
- (c) a nucleic acid sequence substantially identical to (a) or (b).

33. The method of claim 24, wherein said non-cardiomyocyte is a stem or progenitor cell.

5 34. The method of claim 33, wherein said stem or progenitor cell is selected from the group consisting of embryonic and adult stem or progenitor cells.

10 35. The method of claim 33, wherein said stem or progenitor cell is selected from the group consisting of circulating and non-circulating stem or progenitor cells.

36. The method of claim 24, wherein said non-cardiomyocyte is autologous to said animal.

15 37. The method of claim 36, said method further comprising obtaining said non-cardiomyocyte from said animal prior to inducing said differentiation.

20 38. The method of claim 24, wherein said non-cardiomyocyte is non-autologous to said animal.

39. The method of claim 38, wherein said non-cardiomyocyte is allogenic to said animal.

25 40. The method of claim 38, wherein said non-cardiomyocyte is xenogenic to said animal.

30 41. The method of claim 24, wherein said cardiomyocyte is characterized by an alteration of a phenotypic feature relative to said non-cardiomyocyte, wherein said phenotypic feature is selected from the group consisting of:

- 35 (a) level of oxytocin receptor (OTR) protein or OTR-encoding nucleic acid;
- (b) level of ANP protein or ANP-encoding nucleic acid;

- (c) level of muscular MHC protein or muscular MHC-encoding nucleic acid;
- (d) level of DHPR-alpha1 protein or DHPR-alpha1-encoding nucleic acid;
- 5 (e) level of sarcomeric marker proteins;
- (f) level of ion channels;
- (g) mitochondrial dye retention;
- (h) appearance of rhythmic beats; and
- (i) chronotropic responses.

10


42. Use of an agent capable of stimulating OTR activity for treating a disease characterized by cardiomyocyte loss or deficiency in an animal.

15

43. The use of claim 42, wherein said agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).

20

44. The use of claim 43, wherein said oxytocin or functional derivative thereof has the structure:



25

wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.

30

45. The use of claim 42, wherein said animal is a mammal.

46. The use of claim 42, wherein said animal is a human.


47. The use of claim 42, where said disease is selected from the group consisting of cardiac congenital dysfunctions,

aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial ischemia.

48. A commercial package comprising an agent capable of stimulating OTR activity together with instructions for treating a disease characterized by cardiomyocyte loss or deficiency in an animal.

49. The commercial package of claim 48, wherein said agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).

50. The commercial package of claim 49, wherein said oxytocin or functional derivative thereof has the structure:



wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.

51. The commercial package of claim 48, wherein said animal is a mammal.

52. The commercial package of claim 48, wherein said animal is a human.

53. The commercial package of claim 48, where said disease is selected from the group consisting of cardiac congenital dysfunctions, aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial ischemia.


54. A commercial package comprising an agent capable of stimulating OTR activity together with instructions for inducing differentiation of a non-cardiomyocyte to a cardiomyocyte.

5

55. The commercial package of claim 54, wherein said agent is selected from the group consisting of oxytocin or a functional derivative thereof, retinoic acid and triiodothyronine (T<sub>3</sub>).

10

56. The commercial package of claim 55, wherein said oxytocin or functional derivative thereof has the structure:



wherein R is selected from the group consisting of OH, NH<sub>2</sub>, Gly, Gly-Lys and Gly-Lys-Arg.

20 57. A commercial package comprising a culture medium comprising oxytocin or a functional derivative thereof; together with instructions for culturing a non-cardiomyocyte in said culture medium thereby to induce differentiation of said non-cardiomyocyte into a  
25 cardiomyocyte.

58. The commercial package of claim 57, wherein said non-cardiomyocyte is a mammalian non-cardiomyocyte.

30 59. The commercial package of claim 57, wherein said non-cardiomyocyte is a human non-cardiomyocyte.

60. The commercial package of claim 57, wherein said non-cardiomyocyte is a stem or progenitor cell.

35

61. The commercial package of claim 57, wherein said oxytocin or functional derivative thereof is present in said medium at a concentration from about  $10^{-10}$  M to about  $10^{-4}$  M.

5

62. The commercial package of claim 61, wherein said oxytocin or functional derivative thereof is present in said medium at a concentration from about  $10^{-9}$  M to about  $10^{-6}$  M.

10

63. The commercial package of claim 62, wherein said oxytocin or a functional derivative thereof is present in said medium at a concentration from about  $10^{-8}$  M to about  $10^{-7}$  M.

15

64. A composition for treatment of a disease characterized by cardiomyocyte loss or deficiency comprising oxytocin or a functional derivative thereof in and a pharmaceutically acceptable carrier.

20

65. The composition of claim 64, wherein said disease is selected from the group consisting of cardiac congenital dysfunctions, aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial ischemia.

25

66. A method of identifying or characterizing a compound for inducing differentiation of a non-cardiomyocyte cell into a cardiomyocyte, said method comprising:

30

(d) contacting a test compound with a cell comprising an oxytocin receptor (OTR) or an OTR-encoding nucleic acid; and

(e) determining whether OTR activity or expression is increased in the presence of the test compound, said increase in OTR activity or expression being an

35



indication that the test compound may be used for inducing differentiation of a non-cardiomyocyte into a cardiomyocyte.

- 5 67. A method of identifying or characterizing a compound for treatment of a disease characterized by cardiomyocyte loss or deficiency, said method comprising:
- 10 (a) contacting a test compound with a cell comprising an oxytocin receptor (OTR) or an OTR-encoding nucleic acid; and
- 15 (b) determining whether OTR activity or expression is increased in the presence of the test compound, said increase in OTR activity or expression being an indication that the test compound may be used for for treatment of a disease characterized by cardiomyocyte loss or deficiency.
- 20 68. The method of claim 67, wherein said disease is selected from the group consisting of cardiac congenital dysfunctions, aging-related heart pathologies, heart infarction, congestive heart failure and acute myocardial ischemia.
- 25 69. A method of identifying a cell capable of differentiation to a cardiomyocyte, said method comprising determining whether OTR activity or expression is present in said cell, said presence being an indication that said cell is capable of differentiation to a cardiomyocyte.

1/15




FIG. 1

2/15



FIG. 2A




FIG. 2B

3/15





FIG. 3A

4/15

**FIG. 3B**





FIG. 3C

6/15




FIG. 4A




FIG. 4B




FIG. 4C

7/15




FIG. 5A




FIG. 5B



8/15




FIG. 6

9/15




FIG. 7A






FIG. 7B

10/15



RA  $10^{-8}$  M + OTA  $10^{-7}$  M

**FIG. 8B**



RA  $10^{-8}$  M

**FIG. 8A**

11/15




FIG. 9

12/15




FIG. 10B




FIG. 10D




FIG. 10A




FIG. 10C

13/15

Human oxytocin DNA and polypeptide sequences (Genbank  
accession NM\_000915)

Human OT DNA (SEQ ID NO:3):

```
1 accagtcacg gaccctggac ccagcgcacc cgcaccatgg ccggccccag cctcgcttgc
61 tgtctgctcg gcttcctggc gctgacctcc gcttgctaca tccagaactg ccccctggga
121 ggcaagaggg ccgcgcggga cctcgacgtg cgcaagtgcc tcccctgcgg ccccgggggc
181 aaaggccgct gcttcgggac caatatctgc tgcgcggaag agctgggctg cttcgtgggc
241 accgccgaag cgctgcgctg ccaggaggag aactacctgc cgtcgccctg ccagtcgggc
301 cagaaggcgt gcgggagcgg gggccgctgc gcggtcttgg gcctctgctg cagcccggac
361 ggctgccacg ccgacctgac ctgcgacgcg gaagccacct tctcccagcg ctgaaacttg
421 atggctccga acacctcga agcgcgccac tcgcttcccc catagccacc ccagaaatgg
481 tgaaaataaa ataaagcagg tttttctcct ct
```

Encoded polypeptide (SEQ ID NO:4):

MAGPSLACLLGLLALTSACYIQNCPLG GKRAAPDL DV RKCLPCGPGGKGR CFGPNICCAEELGCFVGTAEALR  
CQEENYLPSPCQSGQKACGSGGRCAVLGLCCSPDGCHADPACDAEATFSQR

human OT-encoding region (SEQ ID NO:5):

tgctaca tccagaactg ccccctggga

human OT peptide sequence (SEQ ID NO:6):

CYIQNCPLG

**FIG. 11**

14/15

Human oxytocin receptor DNA and polypeptide  
sequences (Genbank accession NM\_000916)

DNA sequence (SEQ ID NO:7):

```

1  tgttaaggct  ctgggaccaa  cgctgggcga  accagctccg  ctccggaggg  gtctgcgcgg
61  ctggcctcgc  ccgcccccta  gcggaccctg  gcgatatgtc  agcctcagcc  ccaggcacag
121  cgccgcatcc  agacgccgtc  cgcgcgcgca  gcctgggagg  cgctcctcgc  tcgcctcctg
181  taccatcca  gcgaccagcc  aggctgcggc  gaggggattc  caaccgaggc  tccagtgaga
241  gacctcagct  tagcatcaca  ttaggtgcag  ccggcaggcc  atcccaactc  gggccgggag
301  cgcacgcgtc  actggggccg  tcatcgccg  tgcaacttcc  ccggggggag  tcaactttag
361  gttcgctcgc  ggactcgggt  cagtggaaac  cgctgaacat  cccgaggaac  tggcacgctg
421  ggggctctgg  gcttgtggcc  ggtagaggat  tcccgtcat  ttgcagtggc  tcagaggagg
481  gtggaccag  cagatccgtc  cgtggagtct  ccaggagtgg  agccccgggc  gccctacac
541  cctccgacac  gccggatccg  gccagccgc  gccaaagccg  aaagggctcg  aaggccgggg
601  cgcaccgtg  ccgccagggt  catggagggc  gcgctcgcag  ccaactggag  cccgaggcca
661  gccaacgcca  gcgccgcgcc  gccggggccc  gagggcaacc  gcaccgccgg  acccccgcgg
721  cgcaacgagg  ccctggcgcg  cgtggagggt  gcggtgctgt  gtctcatcct  gctcctggcg
781  ctgagcggga  acgcgtgtgt  gctgctggcg  ctgcgcacca  caccgcagaa  gcactcgcgc
841  ctcttcttct  tcatgaagca  cctaagcatc  gccgacctgg  tgggtggcagt  gtttcagggt
901  ctgccgcagt  tgctgtggga  catcaccttc  cgcttctacg  ggcccagact  gctgtgccgc
961  ctggtcaagt  acttgcaagg  ggtgggcatg  ttgcctcca  cctacctgct  gctgctcatg
1021  tccctggacc  gctgcctggc  catctgccag  ccgctgcgct  cgctgcgcgc  ccgcaccgac
1081  gccttgccag  tgctcgccac  gtggctcggc  tgcttggtgg  ccagcgcgcc  gcaggtgcac
1141  atcttctctc  tgccgcagggt  ggctgacggc  gtcttcgact  gctgggcccgt  cttcatccag
1201  ccctggggac  ccaaggccta  catcacatgg  atcacgctag  ctgtctacat  cgtgccgggtc
1261  atcgtgctcg  ctacctgcta  cgcccttatt  agcttcaaga  tctggcagaa  cttgcggctc
1321  aagaccgctg  cagcggcggc  ggccgaggcg  ccagagggcg  cggcggctgg  cgatgggggg
1381  cgctggccc  tggcgcgtgt  cagcagcgtc  aagctcatct  ccaaggccaa  gatccgcacg
1441  gtcaagatga  ctttcatcat  cgtgctggcc  ttcatcgtgt  gctggacgcc  tttcttcttc
1501  gtgcagatgt  ggagcgtctg  ggatgccaac  gcgccaagg  aagcctcggc  cttcatcatc
1561  gtcagtctcc  tggccagcct  caacagctgc  tgcaaccct  ggatctacat  gctgttcacg
1621  ggccacctct  tccacgaact  cgtgcagcgc  ttctgtgct  gctccgccag  ctacctgaag
1681  ggcagacgcc  tgggagagac  gagtgccagc  aaaaagagca  actcgtcctc  cttgtcctg
1741  agccatcgca  gctccagcca  gaggaagctg  tcccagccat  ccacggcggt  acccaccagc
1801  cagggccagg  gctgcagcct  gaggtcagg  ctgtgctggc  ataagtgtct  tgctcctagg
1861  tgatggcgta  tgtttggtga  taaggtacct  atcagtttgt  atccctcccc  tccttgggggt
1921  ggcttcagt  ggggtggag  tggcctccat  gatggaagat  gataggggac  tcagccatca
1981  gacaacaccc  tggcctccta  cacgtacttc  taccaccctg  aaccactgc  tgccctgggc
2041  agtgagtggc  ttgttttttc  tctggacttc  gtaatttcac  tccagtatat  ttttacttct
2101  tcattctggg  atattgtgaa  aagcggtaaa  tataggattg  gtgaccaatt  gggtcaggaa
2161  gtccagtgtt  ctggacttgg  ggtaagcagt  ggggttggga  cctcagatgg  gaaggggtgt
2221  gctaagatcc  tcctgacctc  aaagtgtatt  tgcttttaag  cgaacaaatg  ctggggtcct
2281  tggggaccag  cttgtcagag  ggtagcccta  agagaagggg  attaccttgt  aagaccatct
2341  ggcgagtggt  acctattaga  acttgggtta  aaaaatgtta  agaagctaata  gtttaagaag
2401  catttgggaa  agaaaaagaa  ataaatgtat  ccagatagga  aaagaagaag  taaaactatt
2461  tgcagatgac  acagttttgt  atatagaaaa  tcctaaggaa  ctcacacaca  cacacacaca
2521  cacacagcca  cacagctatt  agaactaata  agcaagttcc  gcaaggtttc  aagatacaag
2581  atcaatatac  aaaaatgaat  tgtatttctt  tatactagca  acaacaata  tgaaaacgaa
2641  gttaaataat  tccatttata  ataccatcag  aaagaataaa  ataggaatca  acttaacaaa
2701  acaagtgcaa  gactgaaaac  tacaaaattg  gaaagaaatt  aaagaaggct  taaataaatg
2761  gaaagacatc  ctgtgttcac  ggatcagact  tagtattgtt  aagatggcaa  tactactcta
2821  actgacatgc  agattcagt  caatccttat  gaaaatcata  gctggctttt  ttacagaaat
2881  tgataagcta  gtcccaaat  tcataaagaa  atgcaaggga  cccagatata  caaataagcc
2941  ttgaaaaaga  acaaagttgg  tggattcaca  cttcctgatt  tcataattta  cgataaagggt
3001  aatcagctca  gtgtgttact  ggtttaagga  tagacatacg  gagcagaata  aagagtacag

```

FIG. 12

15/15

```

3061 atatgaacac ttatacttac ggtcaattga tttttgacaa gggtcccaag acaattcaat
3121 agagaaaagga gagtcttttc aacaaatggc accgagacaa tgatatgcaa gtgcaaaaaga
3181 atgaggttgg acctttactc acactatgtg caaaaatcaa ctcaaaacgc atccaagatc
3241 taaatataag agctgaaact ataaaaatctt agaaaagaaac ataggcatag atctttgtaa
3301 ccttgaatta ggcagtggtt tcttagatat gataccaaaag acacaagcaa ccaatggaaa
3361 aataggtaaa ttggacttaa tcaagatttg aagcttttgt gattgaaaag accctatcaa
3421 gaaggtgaaa agataacctg cagaatggga gaaaatattt gcgagtcata tatatgataa
3481 ggggcttgta tctggaatat ataaataact cttataacac aacaataagg agaaaaataa
3541 atcaatttaa aaaatgggct aacggtttga atagacattt ctccaaagaa gatatgcaaa
3601 tggctactaa gcacatgaaa aatactcaac attattattc attagggaaa tgcaagtcaa
3661 aatcacaaatg agattccagt ttacaatcac taggatggct acaataaaaa gatggacaag
3721 aacgagtgtc ggtgaggatg tagagaaact ggtagaaatt taaattgttg gtgggaatgt
3781 aaatggtgca cctgctttga aaaacagttt ggcagtacct caaaaagtta accgtagagt
3841 gaccatatga cccaggaatg ccactcctag gtatttacc cagagaaatg aaaacgtaca
3901 tacacacaaa aacttgta ccaatgttca tagcaacatt atttgtaata gccaaaaagt
3961 ggaaacaacc caaatgtcta ccaactgatg aatgggaaat aaaatgtggt ctgtccacgc
4021 aatggaacat tattagactc taaaaagaaa tgaagtactc acacatgcca caacatggat
4081 gagccttgaa aacttgctaa gtgaaagaag ccaggtgcaa aagcccacat attgtctgac
4141 tgcattgaaa tgcaatgtct aaaatggacg aatctatata gagtgaatat agattagcgt
4201 ttgccagggc ctggaggctg tgagagatga ggcataacta ctaaggggtt ggggtttctt
4261 tttcgggtga tgaaaatgtt cgaaattagt ggtgattgtg cacgattttg agaatgtact
4321 aaaaaccaat gaactttaaa aaataaaaaa aaacaaa

```

Polypeptide sequence (SEQ ID NO:8):

```

MEGALAANWSAEAAANASAAPPGAEGNRTAGPPRRNEALARVEVAVLCLILLLALSGNACVLLALRTRQKHSRLFFF
MKHLSIADLVVAVFQVLPQLLWDITFRFYGPDLLCRLVKYLQVVGMFASYLLLLMSLDRCLAICQPLRSLRRRTDR
LAVLATWLGCLVASAPQVHIFSLREVADGVFDCWAVFIQPWGPKAYITWITLAVYIVPVIVLATCYGLISFKIWQNL
RLKTAATAAAAEAEPEGAAAGDGGRVALARVSSVKLISKAKIRTVKMTFIIIVLAFIVCWTPFFVFQMWVWDANAPKEA
SAFIIIVMLLASLNSCCNPWIYMLFTGHLFHELVQRFLCCSASYLKGRRLGETSASKKSNSSSFVLSHRSSSQRSQSQ
PSTA

```

**FIG. 12**  
(CONTINUED)



1/10  
SEQUENCE LISTING

<110> UNIVERSITE DU QUEBEC A MONTREAL  
CENTRE HOSPITALIER DE L'UNIVERSITE DE MONTREAL  
PAQUIN, Joanne  
GUTKOWSKA, Jolanta  
DANALACHE, Bogdan A.  
JANKOWSKI, Marek

<120> OXYTOCIN AS CARDIOMYOGENESIS INDUCER AND USES THEREOF

<130> 86197-10

<150> CA 2,391,118

<151> 2002-06-21

<160> 15

<170> PatentIn version 3.1

<210> 1 .

<211> 9

<212> PRT

<213> Homo sapiens

<220>

<221> DISULFID

<222> (1)..(6)

<223>

<220>

<221> MOD\_RES

<222> (9)..(9)

<223> C-terminal Gly is amidated

<400> 1

Cys Tyr Ile Gln Asn Cys Pro Leu Gly  
1 5

<210> 2

<211> 9

<212> PRT

<213> Homo sapiens

<220>

<221> DISULFID

<222> (1)..(6)

<223>

<400> 2

Cys Tyr Ile Gln Asn Cys Pro Leu Gly  
1 5

2/10

```
<210> 3
<211> 512
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (37) .. (414)
<223>
```

[illegible]

```
<210> 4
<211> 125
<212> PRT
<213> Homo sapiens
```

3/10

&lt;400&gt; 4

Met Ala Gly Pro Ser Leu Ala Cys Cys Leu Leu Gly Leu Leu Ala Leu  
 1 5 10 15

Thr Ser Ala Cys Tyr Ile Gln Asn Cys Pro Leu Gly Gly Lys Arg Ala  
 20 25 30

Ala Pro Asp Leu Asp Val Arg Lys Cys Leu Pro Cys Gly Pro Gly Gly  
 35 40 45

Lys Gly Arg Cys Phe Gly Pro Asn Ile Cys Cys Ala Glu Glu Leu Gly  
 50 55 60

Cys Phe Val Gly Thr Ala Glu Ala Leu Arg Cys Gln Glu Glu Asn Tyr  
 65 70 75 80

Leu Pro Ser Pro Cys Gln Ser Gly Gln Lys Ala Cys Gly Ser Gly Gly  
 85 90 95

Arg Cys Ala Val Leu Gly Leu Cys Cys Ser Pro Asp Gly Cys His Ala  
 100 105 110

Asp Pro Ala Cys Asp Ala Glu Ala Thr Phe Ser Gln Arg  
 115 120 125

<210> 5  
 <211> 27  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(27)  
 <223>

<400> 5  
 tgc tac atc cag aac tgc ccc ctg gga  
 Cys Tyr Ile Gln Asn Cys Pro Leu Gly  
 1 5

27

<210> 6  
 <211> 9  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 6

Cys Tyr Ile Gln Asn Cys Pro Leu Gly  
 1 5

```
<210> 7
<211> 4357
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (622)..(1791)
<223>
```

[illegible]

5/10

ggg ccc gac ctg ctg tgc cgc ctg gtc aag tac ttg cag gtg gtg ggc Gly Pro Asp Leu Leu Cys Arg Leu Val Lys Tyr Leu Gln Val Val Gly 110 115 120	987
atg ttc gcc tcc acc tac ctg ctg ctg ctc atg tcc ctg gac cgc tgc Met Phe Ala Ser Thr Tyr Leu Leu Leu Leu Met Ser Leu Asp Arg Cys 125 130 135	1035
ctg gcc atc tgc cag ccg ctg cgc tcg ctg cgc cgc cgc acc gac cgc Leu Ala Ile Cys Gln Pro Leu Arg Ser Leu Arg Arg Thr Asp Arg 140 145 150	1083
ctg gca gtg ctc gcc acg tgg ctc ggc tgc ctg gtg gcc agc gcg ccg Leu Ala Val Leu Ala Thr Trp Leu Gly Cys Leu Val Ala Ser Ala Pro 155 160 165 170	1131
cag gtg cac atc ttc tct ctg cgc gag gtg gct gac ggc gtc ttc gac Gln Val His Ile Phe Ser Leu Arg Glu Val Ala Asp Gly Val Phe Asp 175 180 185	1179
tgc tgg gcc gtc ttc atc cag ccc tgg gga ccc aag gcc tac atc aca Cys Trp Ala Val Phe Ile Gln Pro Trp Gly Pro Lys Ala Tyr Ile Thr 190 195 200	1227
tgg atc acg cta gct gtc tac atc gtg ccg gtc atc gtg ctc gct acc Trp Ile Thr Leu Ala Val Tyr Ile Val Pro Val Ile Val Leu Ala Thr 205 210 215	1275
tgc tac ggc ctt atc agc ttc aag atc tgg cag aac ttg cgg ctc aag Cys Tyr Gly Leu Ile Ser Phe Lys Ile Trp Gln Asn Leu Arg Leu Lys 220 225 230	1323
acc gct gca gcg gcg gcg gcc gag gcg cca gag ggc gcg gcg gct ggc Thr Ala Ala Ala Ala Ala Ala Glu Ala Pro Glu Gly Ala Ala Ala Gly 235 240 245 250	1371
gat ggg ggg cgc gtg gcc ctg gcg cgt gtc agc agc gtc aag ctc atc Asp Gly Gly Arg Val Ala Leu Ala Arg Val Ser Ser Val Lys Leu Ile 255 260 265	1419
tcc aag gcc aag atc cgc acg gtc aag atg act ttc atc atc gtg ctg Ser Lys Ala Lys Ile Arg Thr Val Lys Met Thr Phe Ile Ile Val Leu 270 275 280	1467
gcc ttc atc gtg tgc tgg acg cct ttc ttc ttc gtg cag atg tgg agc Ala Phe Ile Val Cys Trp Thr Pro Phe Phe Phe Val Gln Met Trp Ser 285 290 295	1515
gtc tgg gat gcc aac gcg ccc aag gaa gcc tcg gcc ttc atc atc gtc Val Trp Asp Ala Asn Ala Pro Lys Glu Ala Ser Ala Phe Ile Ile Val 300 305 310	1563
atg ctc ctg gcc agc ctc aac agc tgc tgc aac ccc tgg atc tac atg Met Leu Leu Ala Ser Leu Asn Ser Cys Cys Asn Pro Trp Ile Tyr Met 315 320 325 330	1611
ctg ttc acg ggc cac ctc ttc cac gaa ctc gtg cag cgc ttc ctg tgc Leu Phe Thr Gly His Leu Phe His Glu Leu Val Gln Arg Phe Leu Cys 335 340 345	1659

6/10

tgc tcc gcc agc tac ctg aag ggc aga cgc ctg gga gag acg agt gcc	1707
Cys Ser Ala Ser Tyr Leu Lys Gly Arg Arg Leu Gly Glu Thr Ser Ala	
350 355 360	
agc aaa aag agc aac tcg tcc tcc ttt gtc ctg agc cat cgc agc tcc	1755
Ser Lys Lys Ser Asn Ser Ser Ser Phe Val Leu Ser His Arg Ser Ser	
365 370 375	
agc cag agg agc tgc tcc cag cca tcc acg gcg tga cccaccagcc	1801
Ser Gln Arg Ser Cys Ser Gln Pro Ser Thr Ala	
380 385	
agggccaggg ctgcagcctg aggctcaggc tgtgctggca taagtgtctt gtccttaggt	1861
gatggcgtat gtttgtgtat aaggtaacct tcagtttgta tccctcccct ccttgggggtg	1921
gcttcagtgg ggtggagagt ggctccatg atggaagatg ataggggact cagccatcag	1981
acaacaccct ggctcctac acgtacttct accaccctga acccactgct gccctgggca	2041
gtgagtggct tgttttttct cctggacttg taatttctact ccagtatatt tttacttctt	2101
cattctggga tattgtgaaa agcggtaa ataggattgg tgaccaattg ggtcaggaag	2161
tccagtgttc tggacttggg gtaagcagtg gggttgggac ctcagatggg aagggtggtg	2221
ctaagatcct cctgacctca aagtgtattt gcctttaagc gaacaaatgc tggggtcctt	2281
ggggaccagc ttgtcagagg gtagccctaa gagaagggga ttaccttgta agaccatctg	2341
gcgcagtgga cctattagaa cttgggttaa aaatgtttta gaagctaattg tttaagaagc	2401
atttgggaaa gaaaaagaaa taaatgtatc cagataggaa aagaagaagt aaaactattt	2461
gcagatgaca cagttttgta tatagaaaat cctaaggaa tcacacacac acacacacac	2521
acacacgcac acagctatta gaactaataa gcaagttccg caaggtttca agatacaaga	2581
tcaatataca aaaatgaatt gtatttcttt atactagcaa caaacaatat gaaaacgaag	2641
ttaaataatt ccatttataa taccatcaga aagaataaaa taggaatcaa cttaacaaaa	2701
caagtgcaag actgaaaact acaaaattgg aaagaaatta aagaaggctt aaataaatgg	2761
aaagacatcc tgtgttcctg gatcagactt agtattgtta agatggcaat actatcctaa	2821
ctgacatgca gattcagtgc aatccttatg aaaatcatag ctggcttttt tacagaaatt	2881
gataagctag tcccaaaatt cataaagaaa tgcaagggac ccagatatcc aaataagcct	2941
tgaaaaagaa caaagttggg ggattcacac ttcttgattt cataatttac gataaaggta	3001
atcagctcag tgtgttactg gtttaaggat agacatacgg agcagaataa agagtacaga	3061
tatgaacact tatacttacg gtcaattgat ttttgacaag gttcccaaga caattcaata	3121
gagaaaggag agtcttttca acaaattggc ccgagacaat gatatgcaag tgcaaaagaa	3181
tgaggttggg cttttactca cactatgtgc aaaaatcaac tcaaaacgca tccaagatct	3241

7/10

```

aaatataaga gctgaaacta taaaatctta gaaagaaaca taggcataga tctttgtaac 3301
cttgaattag gcagtggttt cttagatatg ataccaaaga cacaagcaac caatggaaaa 3361
ataggtaaat tggacttaat caagatttga agcttttgtg attgaaaaga ccctatcaag 3421
aaggtgaaaa gataacctgc agaatgggag aaaatatttg cgagtcatat atatgataag 3481
gggcttgat ctggaatata taaataactc ttataacaca acaataagga gaaaaataaa 3541
tcaatttaaa aaatgggcta acggtttgaa tagacatttc tccaaagaag atatgcaaat 3601
ggctactaag cacatgaaaa atactcaaca ttattattca ttagggaaat gcaagtcaaa 3661
atcacaatga gattccagtt tacaatcact aggatggcta caataaaaag atggacaaga 3721
acgagtgtcg gtgaggatgt agagaaaactg gtagaaattt aaattgttg tgggaatgta 3781
aatggtgcac ctgctttgaa aaacagtttg gcagtacctc aaaaagttaa acgtagagtg 3841
accatatgac ccaggaatgc cactcctagg tatttaccga agagaaatga aaacgtacat 3901
acacacaaaa acttgtaacac caatgttcat agcaacatta ttgtaatag ccaaaaagtg 3961
gaaacaaccc aaatgtctac caactgatga atgggaaata aaatgtggtc tgtccacgca 4021
atggaacatt attagactct aaaaagaaat gaagtactca cacatgccac aacatggatg 4081
agccttgaaa acttgctaag tgaaagaagc caggtgcaaa agccacata ttgtctgact 4141
gcattgaaat gcaatgtcta aaatggacga atctatatag agtgaatata gattagcggt 4201
tgccagggcc tggaggctgt gagagatgag gcatgactac taagggttg gggtttcttt 4261
ttcgggtgat gaaaatgttc gaaattagtg gtgattgtgc acgattttga gaatgtacta 4321
aaaaccaatg aactttaaaa aataaaaata aacaaa 4357

```

<210> 8  
 <211> 389  
 <212> PRT  
 <213> Homo sapiens

<400> 8

```

Met Glu Gly Ala Leu Ala Ala Asn Trp Ser Ala Glu Ala Ala Asn Ala
1           5           10           15

```

```

Ser Ala Ala Pro Pro Gly Ala Glu Gly Asn Arg Thr Ala Gly Pro Pro
20           25           30

```

```

Arg Arg Asn Glu Ala Leu Ala Arg Val Glu Val Ala Val Leu Cys Leu
35           40           45

```

```

Ile Leu Leu Leu Ala Leu Ser Gly Asn Ala Cys Val Leu Leu Ala Leu
50           55           60

```

8/10

Arg Thr Thr Arg Gln Lys His Ser Arg Leu Phe Phe Phe Met Lys His  
65 70 75 80

Leu Ser Ile Ala Asp Leu Val Val Ala Val Phe Gln Val Leu Pro Gln  
85 90 95

Leu Leu Trp Asp Ile Thr Phe Arg Phe Tyr Gly Pro Asp Leu Leu Cys  
100 105 110

Arg Leu Val Lys Tyr Leu Gln Val Val Gly Met Phe Ala Ser Thr Tyr  
115 120 125

Leu Leu Leu Leu Met Ser Leu Asp Arg Cys Leu Ala Ile Cys Gln Pro  
130 135 140

Leu Arg Ser Leu Arg Arg Arg Thr Asp Arg Leu Ala Val Leu Ala Thr  
145 150 155 160

Trp Leu Gly Cys Leu Val Ala Ser Ala Pro Gln Val His Ile Phe Ser  
165 170 175

Leu Arg Glu Val Ala Asp Gly Val Phe Asp Cys Trp Ala Val Phe Ile  
180 185 190

Gln Pro Trp Gly Pro Lys Ala Tyr Ile Thr Trp Ile Thr Leu Ala Val  
195 200 205

Tyr Ile Val Pro Val Ile Val Leu Ala Thr Cys Tyr Gly Leu Ile Ser  
210 215 220

Phe Lys Ile Trp Gln Asn Leu Arg Leu Lys Thr Ala Ala Ala Ala Ala  
225 230 235 240

Ala Glu Ala Pro Glu Gly Ala Ala Ala Gly Asp Gly Gly Arg Val Ala  
245 250 255

Leu Ala Arg Val Ser Ser Val Lys Leu Ile Ser Lys Ala Lys Ile Arg  
260 265 270

Thr Val Lys Met Thr Phe Ile Ile Val Leu Ala Phe Ile Val Cys Trp  
275 280 285

Thr Pro Phe Phe Phe Val Gln Met Trp Ser Val Trp Asp Ala Asn Ala  
290 295 300



9/10

Pro Lys Glu Ala Ser Ala Phe Ile Ile Val Met Leu Leu Ala Ser Leu  
 305 310 315 320

Asn Ser Cys Cys Asn Pro Trp Ile Tyr Met Leu Phe Thr Gly His Leu  
 325 330 335

Phe His Glu Leu Val Gln Arg Phe Leu Cys Cys Ser Ala Ser Tyr Leu  
 340 345 350

Lys Gly Arg Arg Leu Gly Glu Thr Ser Ala Ser Lys Lys Ser Asn Ser  
 355 360 365

Ser Ser Phe Val Leu Ser His Arg Ser Ser Ser Gln Arg Ser Cys Ser  
 370 375 380

Gln Pro Ser Thr Ala  
 385

<210> 9  
 <211> 22  
 <212> DNA  
 <213> Artificial

<220>  
 <223> synthetic oligonucleotide

<400> 9  
 aagatgacct tcattcattgt tc

22

<210> 10  
 <211> 23  
 <212> DNA  
 <213> Artificial

<220>  
 <223> synthetic oligonucleotide

<400> 10  
 cgactcagga cgaaggtgga gga

23

<210> 11  
 <211> 24  
 <212> DNA  
 <213> Artificial

<220>  
 <223> synthetic oligonucleotide

<400> 11  
 gtcaatccta cccccgaagc agct

24

10/10

<210> 12  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> synthetic oligonucleotide

<400> 12  
cagcatgggc tccttctcca 20

<210> 13  
<211> 23  
<212> DNA  
<213> Artificial

<220>  
<223> synthetic oligonucleotide

<400> 13  
cagtgatggc atccactgtg gtc 23

<210> 14  
<211> 27  
<212> DNA  
<213> Artificial

<220>  
<223> synthetic oligonucleotide

<400> 14  
aaggtcgggtg tcaaccatt tggccgt 27

<210> 15  
<211> 10  
<212> PRT  
<213> Homo sapiens

<220>  
<221> DISULFID  
<222> (1)..(6)  
<223>

<400> 15

Cys Tyr Ile Gln Asn Cys Pro Leu Gly Gly  
1 5

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number  
**WO 2004/000993 A3**

- (51) International Patent Classification<sup>7</sup>: **C12N 5/06**, 5/08, A61K 38/11
- (21) International Application Number: **PCT/CA2003/000897**
- (22) International Filing Date: 13 June 2003 (13.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
2,391,118 21 June 2002 (21.06.2002) CA
- (71) Applicants (*for all designated States except US*): **UNIVERSITE DU QUEBEC A MONTREAL** [CA/CA]; Succursale Centre-Ville, Case Postale 8888, Montreal, Quebec H3C 3P8 (CA). **CENTRE HOSPITALIER DE L'UNIVERSITE DE MONTREAL** [CA/CA]; Centre de Recherche Hotel Dieu, Pavillon Masson, 3850 Rue St-Urbain, Montreal, Quebec H2W 1T7 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **PAQUIN, Joanne** [CA/CA]; 10151 Georges-Baril, Montréal, Québec H2C 2M9 (CA). **GUTKOWSKA, Jolanta** [CA/CA]; 139 Avenue Appin, Ville Mont-Royal, Quebec H3P 1V6 (CA). **DANALACHE, Bogdan, A.** [RO/CA]; 4555 St. Kevin, App. 306, Montreal, Quebec H3T 1J1 (CA). **JANKOWSKI, Marek** [CA/CA]; 5155 West Broadway, Montreal, Quebec H4V 2A1 (CA).
- (74) Agents: **SHAHINIAN, Serge, S. et al.; SMART & BIGGAR**, 1000 de la Gauchetière Street West, Suite 3400, Montréal, Québec H3B 4W5 (CA).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— *with international search report*
- (88) Date of publication of the international search report:  
29 April 2004
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: OXYTOCIN AS CARDIOMYOGENESIS INDUCER AND USES THEREOF

(57) Abstract: The invention relates to oxytocin and oxytocin-related compounds and functional derivatives thereof, and uses thereof to induce differentiation of a non-cardiomyocyte (e.g. a stem/progenitor cell) to a cardiomyocyte. The invention further relates to the methods of prevention or treatment of conditions characterized by or associated with a cardiomyocyte loss or deficiency, by administration of oxytocin or an oxytocin-related compound or a functional derivative thereof to a subject, or via the administration/transplantation of a cell differentiated ex vivo by a method of the invention. The invention further relates to methods, uses, commercial packages and culture media relating to such differentiation and prevention/treatment.

WO 2004/000993 A3

Internal Application No  
PCT/CA 03/00897

Internal Application No  
PCT/CA 03/00897

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N5/06 C12N5/08 A61K38/11

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N5/06 C12N5/08 A61K38/11

### B. FIELDS SEARCHED

TPC 7 C12N A61K C07K

EPO-Internal, WPI Data, BIOSIS

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

48-65

—/—

☒ Patent family members are listed in annex.

\*& document member of the same patent family

12/12/2003

**Teyssier, B**

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/CA 03/00897

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 01 48149 A (SAKURADA KAZUHIRO ;HATA JUN ICHI (JP); OGAWA SATOSHI (JP); FUKUDA) 5 July 2001 (2001-07-05)</p> <p>-&amp; US 2002/142457 A1 (SAKURADA K ET AL) 3 October 2002 (2002-10-03) examples 3,16</p> <p style="text-align: center;">---</p>	<p>1-3, 7-15, 19-29, 33-43, 45-49, 51-55</p>
X	<p>RODRIGUEZ E R ET AL: "3,5,3'-triiodo-L-thyronine induces cardiac myocyte differentiation but not neuronal differentiation in P19 teratocarcinoma cells in a dose dependent manner" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 205, no. 1, 1994, pages 652-658, XP002263293 ISSN: 0006-291X the whole document</p> <p style="text-align: center;">---</p>	<p>1-3, 7-15, 19-29, 33-43, 45-49, 51-55</p>
A	<p>MCBURNEY M W: "P19 embryonal carcinoma cells" INTERNATIONAL JOURNAL OF DEVELOPMENTAL BIOLOGY, vol. 37, no. 1, 1993, pages 135-140, XP008024916 ISSN: 0214-6282 cited in the application</p> <p style="text-align: center;">---</p>	
P,X	<p>PAQUIN J ET AL: "Oxytocin induces differentiation of P19 embryonic stem cells to cardiomyocytes" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 99, no. 14, 9 July 2002 (2002-07-09), pages 9550-9555, XP002263273 ISSN: 0027-8424 the whole document</p> <p style="text-align: center;">-----</p>	<p>1-69</p>

# INTERNATIONAL SEARCH REPORT

Int... al application No.  
PCT/CA 03/00897

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims 1-9, 11, 12 in part and 13-41 in totality are directed to methods of treatment of animals, including humans. The search has nevertheless been carried out and based on the alleged effects of the composition.
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Present claims 2, 14, 28, 42, 48 and 54 relate to a product and uses thereof, where the product is defined by reference to a desirable property, namely stimulating oxytocin receptor activity. The claims cover all products having this property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search for claims 1, 2, 7-14, 19-28, 33-42, 45-48 and 51-54 has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to oxytocin, oxytocin functional derivatives, nucleic acids encoding oxytocin or functional derivatives thereof, retinoic acid and triiodothyronine, as mentioned in claims 3-6, 15-18, 29-32, 43, 44, 49, 50, 55 and 56.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 03/00897

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0148149	A	05-07-2001	AU 1055201 A	09-07-2001
			AU 2228101 A	09-07-2001
			AU 2695000 A	09-07-2001
			CA 2395950 A1	05-07-2001
			EP 1254952 A1	06-11-2002
			WO 0148149 A1	05-07-2001
			WO 0148150 A1	05-07-2001
			WO 0148151 A1	05-07-2001
			US 2002142457 A1	03-10-2002

---



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**